arXiv:1002.0996v3 [math.GT] 7 Jun 2011

CANNON-THURSTON MAPS FOR KLEINIAN GROUPS

MAHAN MJ

Abstract. We show that Cannon-Thurston maps exist for degenerate free groups without parabolics, i.e. for handlebody groups. Combining these techniques with earlier work proving the existence of Cannon-Thurston maps for surface groups, we show that Cannon-Thurston maps exist for arbitrary finitely generated Kleinian groups, proving a conjecture of McMullen. We also show that point pre-images under Cannon-Thurston maps for degenerate free groups without parabolics correspond to end-points of leaves of an ending lamination in the Masur domain, whenever a point has more than one pre-image. This proves a conjecture of Otal. We also prove a similar result for point pre-images under Cannon-Thurston maps for arbitrary finitely generated Kleinian groups.

Contents

1. Introduction 1
1.1. Relative Hyperbolicity 3
1.2. Cannon-Thurston Maps 5
2. Split Geometry 6
3. Free Groups and Finitely Generated Kleinian Groups 10
3.1. The Masur Domain 11
3.2. Incompressibility of Split Components 11
3.3. Constructing Quasidisks 12
3.4. Reduction Lemma and Admissible Paths 14
3.5. Cannon-Thurston Maps for Free Groups 16
3.6. Finitely Generated Kleinian Groups 18
4. Point pre-images of the Cannon-Thurston Map 20
4.1. EL leaves are CT leaves 20
4.2. CT leaves are EL leaves 22
5. Applications and Extensions 24
5.1. Primitive Stable Representations 24
5.2. Discreteness of Commensurators 25
5.3. Extending the Sullivan-McMullen Dictionary 25
References 27

1. Introduction

In [28] we showed that simply or doubly degenerate surface Kleinian groups without accidental parabolics admit Cannon-Thurston maps, answering affirmatively a question of Cannon and Thurston (Section 6 of [8] [9]). In [29] we had shown that

---

2000 *Mathematics Subject Classification.* 57M50.

point pre-images of the Cannon-Thurston map for simply or doubly degenerate groups without parabolics correspond to endpoints of leaves of ending laminations whenever a point has more than one pre-image. The main aim of this paper is to apply the techniques developed in [28] and the result of [29] to extend these results to arbitrary finitely generated Kleinian groups. This completes the project starting with [28] and proceeding through [29], [10], [11].

The principal new ingredient is a proof of the existence of Cannon-Thurston maps for degenerate handlebody groups without parabolics.[1] The following is the main new ingredient of this paper.

Theorem 4.5 *Let G be a finitely generated free degenerate Kleinian group without parabolics. Let $i : \Gamma_G \to \mathbb{H}^3$ be the natural identification of a Cayley graph of G with the orbit of a point in $\mathbb{H}^3$. Then i extends continuously to a map $\hat{i} : \widehat{\Gamma_G} \to \mathbb{D}^3$. Let ∂i denote the restriction of $\hat{i}$ to the boundary $\partial\Gamma$ of Γ. Then $\partial i(a) = \partial i(b)$ for $a \neq b \in \partial\Gamma$ if and only if a, b are either ideal end-points of a leaf of an ending lamination of G, or ideal boundary points of a complementary ideal polygon.*

A crucial idea in the proof of the existence of the Cannon-Thurston map $\hat{i}$ in Theorem 4.5 goes back to Miyachi [27] in the case of bounded geometry. The proof of Theorem 4.5 generalizes with some modifications to arbitrary finitely generated Kleinian groups.

Theorems 3.19 and 4.6 *Let G be a finitely generated Kleinian group. Let $i : \Gamma_G \to \mathbb{H}^3$ be the natural identification of a Cayley graph of G with the orbit of a point in $\mathbb{H}^3$. Then i extends continuously to a map $\hat{i} : \widehat{\Gamma_G} \to \mathbb{D}^3$, where $\widehat{\Gamma_G}$ denotes the (relative) hyperbolic compactification of Γ_G. Let ∂i denote the restriction of $\hat{i}$ to the boundary $\partial\Gamma_G$ of Γ_G.*

Let E be a degenerate end of $N^h = \mathbb{H}^3/G$ and $\widetilde{E}$ a lift of E to $\widetilde{N^h}$ and let M_{gf} be an augmented Scott core of N^h. Then the ending lamination $\mathcal{L}_E$ for the end E lifts to a lamination on $\widetilde{M_{gf}} \cap \widetilde{E}$. Each such lift $\mathcal{L}$ of the ending lamination of a degenerate end defines a relation $\mathcal{R}_{\mathcal{L}}$ on the (Gromov) Hyperbolic boundary $\partial\widetilde{M_{gf}}$ (equal to the relative hyperbolic boundary $\partial\Gamma_G$ of Γ_G), given by $a\mathcal{R}_{\mathcal{L}}b$ iff a, b are end-points of a leaf of $\mathcal{L}$. Let $\{\mathcal{R}_i\}_i$ be the entire collection of relations on $\partial\widetilde{M_{gf}}$ obtained this way. Let $\mathcal{R}$ be the transitive closure of the union $\bigcup_i \mathcal{R}_i$. Then $\partial i(a) = \partial i(b)$ iff $a\mathcal{R}b$.

Theorem 3.19 gives an affirmative answer to a conjecture of McMullen [22] and Theorem 4.5 gives an affirmative answer to a conjecture of Otal [33].

For ease of exposition, throughout this paper, we shall often first work out the problem for free groups and then indicate the generalization to arbitrary finitely generated Kleinian groups.

Acknowledgments: I am grateful to Jean-Pierre Otal for suggesting the problem of finding point pre-images of the Cannon-Thurston map for handlebodies; and for giving me a copy of his thesis [33], where the structure of Cannon-Thurston maps for handlebody groups is conjectured. I would also like to thank the referee for several suggestions and corrections. This work is partly supported by a CEFIPRA Indo-French Research grant 4301-1.

---

[1] An earlier version of some parts of this paper existed in draft form in an earlier version of [28]. The division of material between [28] and the present paper is in the interests of readability.

1.1. **Relative Hyperbolicity.**

We refer the reader to Farb [12] for terminology and details on relative hyperbolicity and electric geometry.

Definition 1.1. Given a metric space (X, d_X) and a collection $\mathcal{H}$ of subsets, let $\mathcal{E}(X, \mathcal{H}) = X \bigsqcup_{H\in\mathcal{H}} (H \times [0, \frac{1}{2}])$ be the identification space obtained by identifying $(h, 0) \in H \times [0, \frac{1}{2}]$ with $h \in X$. Each $\{h\} \times [0, \frac{1}{2}]$ is declared to be isometric to the interval $[0, \frac{1}{2}]$ and $H \times \{\frac{1}{2}\}$ is equipped with the zero metric. $\mathcal{E}(X, \mathcal{H})$ is given a path pseudo-metric as follows.

Only such paths in $\mathcal{E}(X, \mathcal{H})$ are allowed whose intersection with any $\{h\} \times (0, \frac{1}{2})$ is either all of $\{h\} \times (0, \frac{1}{2})$ or is empty. The distance between two points in $\mathcal{E}(X, \mathcal{H})$ is the infimum of lengths of such allowable paths.

The resulting pseudo-metric space $\mathcal{E}(X, \mathcal{H})$ is the **electric space** associated to X and the collection $\mathcal{H}$.

We shall say that $\mathcal{E}(X, \mathcal{H})$ is constructed from X by **electrocuting** the collection $\mathcal{H}$ and the induced pseudo-metric d_e will be called the **electric metric**.

If $\mathcal{E}(X, \mathcal{H})$ is (Gromov) hyperbolic, we say that X is **weakly hyperbolic** relative to $\mathcal{H}$.

Note that since $\mathcal{E}(X, \mathcal{H}) = X \bigsqcup_{H\in\mathcal{H}} (H \times [0, \frac{1}{2}])$, X can be naturally identified with a subspace of $\mathcal{E}(X, \mathcal{H})$. Paths in (X, d_X) can therefore be regarded as paths in $\mathcal{E}(X, \mathcal{H})$, but are very far from being quasi-isometrically embedded in general.

A collection $\mathcal{H}$ of subsets of (X, d_X) is said to be D-separated if $d_X(H_1, H_2) \geq D$ for all $H_1, H_2 \in \mathcal{H}; H_1 \neq H_2$. D-separatedness is only a technical restriction as the collection $\{H \times \{\frac{1}{2}\} : H \in \mathcal{H}\}$ is 1-separated in $\mathcal{E}(X, \mathcal{H})$.

Definition 1.2. Given a collection $\mathcal{H}$ of C-quasiconvex, D-separated sets in a (Gromov) hyperbolic metric space (X, d_X) we shall say that a geodesic (resp. quasigeodesic) γ is a geodesic (resp. quasigeodesic) **without backtracking** if γ does not return to H after leaving it, for any $H \in \mathcal{H}$.

There is a distinguished collection of 1-separated subsets of $\mathcal{E}(X, \mathcal{H})$ given by $\{H \times \{\frac{1}{2}\} : H \in \mathcal{H}\}$. An electric quasigeodesic **without backtracking** in $\mathcal{E}(X, \mathcal{H})$ is an electric quasigeodesic that does not return to $H \times \{\frac{1}{2}\}$ after leaving it, for any $H \in \mathcal{H}$.

Notation: For any pseudo metric space (Z, ρ) and $A \subset Z$, we shall use the notation $N_R(A, \rho) = \{x \in Z : \rho(x, A) \leq R\}$ as for metric spaces.

Lemma 1.3. *(Lemma 4.5 and Proposition 4.6 of* [12]*; Theorem 5.3 of* [18]*;* [2]*)*
Given δ, C there exists Δ such that if (X, d_X) is a δ-hyperbolic metric space with a collection $\mathcal{H}$ of C-quasiconvex sets. then,
Electric quasi-geodesics electrically track (Gromov) hyperbolic geodesics: *For all $P > 0$, there exists $K > 0$ such that if β is any electric P-quasigeodesic from x to y, and γ is a geodesic in (X, d_X) from x to y, then $\beta \subset N_K(\gamma, d_e)$. Further,*
1) $\gamma \subset N_K((N_0(\beta, d_e)), d_X)$.
2) Relative Hyperbolicity: *X is weakly hyperbolic relative to $\mathcal{H}$. $\mathcal{E}(X, \mathcal{H})$ is Δ-hyperbolic.*

Note that we do not need D-separatedness in the hypothesis of Lemma 1.3.

Let (X, d_X) be a δ-hyperbolic metric space, and $\mathcal{H}$ a family of C-quasiconvex, collection of subsets. Let $\alpha = [a, b]$ be a geodesic in (X, d_X) and β an electric P-quasigeodesic without backtracking in $\mathcal{E}(X, \mathcal{H})$ joining a, b. Order from the left the collection of maximal subsegments of β contained entirely in some $H \times \{\frac{1}{2}\} : H \in \mathcal{H}$. Let $\{[p_i, q_i] \times \{\frac{1}{2}\}(\subset H_i)\}_i$ be the collection of maximal subsegments. Replace, as per this order, each path of the form $\{p_i\} \times [0, \frac{1}{2}] \cup [p_i, q_i] \times \{\frac{1}{2}\} \cup \{q_i\} \times [0, \frac{1}{2}] \subset H_i \times [0, \frac{1}{2}]$ by a geodesic $[p_i, q_i]$ in X. The resulting **connected** path β_q in X is called an *electro-ambient representative* of β in X.

Lemma 1.4. *(See Proposition 4.3 of* [18]*, also see Lemma 3.10 of* [30]*) Given δ, C, P there exists C_3 such that the following holds:*
Let (X, d_X) be a δ-hyperbolic metric space and $\mathcal{H}$ a family of C-quasiconvex subsets. Let (X, d_e) denote the electric space obtained by electrocuting elements of $\mathcal{H}$. Then, if α, β_q denote respectively a (Gromov) hyperbolic geodesic and an electro-ambient P-quasigeodesic with the same end-points, then α lies in a (Gromov hyperbolic d_X-) C_3 neighborhood of β_q.

Two paths β, γ in (X, d_X) with the same endpoints are said to have *similar intersection patterns* with $\mathcal{H}$ if there exists $\epsilon > 0$, depending only on $(X, \mathcal{H})$, such that:

- **Similar Intersection Patterns 1:** If precisely one of $\{\beta, \gamma\}$ meets some $H \in \mathcal{H}$, then the d_X-distance from the entry point to the exit point is at most D.
- **Similar Intersection Patterns 2:** If both $\{\beta, \gamma\}$ meet some $H \in \mathcal{H}$, then the distance from the entry point of β to that of γ is at most D, and similarly for the exit points.

Definition 1.5. [12] Suppose that X is weakly hyperbolic relative to $\mathcal{H}$. Suppose that any two electric quasigeodesics without backtracking and with the same endpoints have similar intersection patterns with respect to the collection $\{H \times \frac{1}{2} : H \in \mathcal{H}\}$. Then $(X, \mathcal{H})$ is said to satisfy **bounded penetration** and X is said to be **strongly hyperbolic** relative to $\mathcal{H}$.

The next condition ensures that $(X, \mathcal{H})$ is **strongly hyperbolic** relative to $\mathcal{H}$.

Definition 1.6. A collection $\mathcal{H}$ of uniformly C-quasiconvex sets in a δ-hyperbolic metric space X is said to be **mutually D-cobounded** if for all $H_i, H_j \in \mathcal{H}$, $\pi_i(H_j)$ has diameter less than D, where π_i denotes a nearest point projection of X onto H_i. A collection is **mutually cobounded** if it is mutually D-cobounded for some D.

Lemma 1.7. [28] *Let X be a hyperbolic metric space and $\mathcal{H}$ a collection of ϵ neighborhoods of mutually cobounded quasiconvex sets; then any electro-ambient quasigeodesic is a quasigeodesic in X.*

Partial Electrocution
Let M be a (not necessarily simply connected) convex hyperbolic 3-manifold with a neighborhood of the cusps excised. Then each boundary component of M is of the form $\sigma \times P$, where P is either an interval or a circle, and σ is a horocycle of some fixed length e_0. In the universal cover $\widetilde{M}$, if we excise (open) horoballs, we are left with a manifold whose boundaries are flat horospheres of the form $\widetilde{\sigma} \times \tilde{P}$. Note that $\tilde{P} = P$ if P is an interval, and $\mathbb{R}$ if P is a circle (the case for a $(Z+Z)$-cusp).

Let Y be a convex simply connected hyperbolic 3-manifold. Let $\mathcal{B}$ denote a collection of horoballs. Let X denote Y minus the interior of the horoballs in $\mathcal{B}$. Let $\mathcal{H}$ denote the collection of boundary horospheres. Then each $H \in \mathcal{H}$ with the induced metric is isometric to a Euclidean product $E^1 \times L$ for an interval $L \subset \mathbb{R}$. Here E^1 denotes Euclidean 1-space. **Partially electrocute** each H by giving it the product of the zero metric with the Euclidean metric, i.e. on E^1 put the zero metric and on L put the Euclidean metric. The resulting space is essentially what one would get (in the spirit of [12]) by gluing to each H the mapping cylinder of the projection of H onto the L-factor. Let d_{pel} denote the partially electrocuted pseudometric on X.

The above construction can be done in the base manifold M itself by equipping the boundary component $\sigma \times P$ with the product of a zero metric in the σ direction and the Euclidean metric in the P-direction.

Lemma 1.8. [28] *(X, d_{pel}) is a (Gromov) hyperbolic metric space.*

1.2. Cannon-Thurston Maps.

Let (X, d_X) and (Y, d_Y) be hyperbolic metric spaces. By adjoining the Gromov boundaries ∂X and ∂Y to X and Y, one obtains their compactifications $\widehat{X}$ and $\widehat{Y}$ respectively.

Let $i : Y \to X$ denote a proper map.

Definition 1.9. *Let X and Y be hyperbolic metric spaces and $i : Y \to X$ be a proper map. A* **Cannon-Thurston map** *$\hat{i}$ from $\widehat{Y}$ to $\widehat{X}$ is a continuous extension of i.*

Lemma 2.1 of [26] below gives a necessary and sufficient condition for the existence of Cannon-Thurston maps.

Lemma 1.10. [26] *A Cannon-Thurston map from $\widehat{Y}$ to $\widehat{X}$ exists iff the following condition is satisfied:*
Given $y_0 \in Y$, there exists a non-negative function $M(N)$, such that $M(N) \to \infty$ as $N \to \infty$ and for all geodesic segments λ lying outside an N-ball around $y_0 \in Y$ any geodesic segment in X joining the end-points of $i(\lambda)$ lies outside the $M(N)$-ball around $i(y_0) \in X$.

We shall now give a criterion for the existence of Cannon-Thurston maps between relatively hyperbolic spaces. Let X and Y be strongly hyperbolic relative to the collections $\mathcal{H}_X$ and $\mathcal{H}_Y$ respectively. Let $i\colon Y \to X$ be a weakly type-preserving proper embedding, i.e. for $H_Y \in \mathcal{H}_Y$ there exists $H_X \in \mathcal{H}_X$ such that $i(H_Y) \subset H_X$ and images of distinct elements of $\mathcal{H}_Y$ lie in distinct elements of $\mathcal{H}_X$.

In the Lemma below, we specialize to the case where X, Y are convex simply connected complete hyperbolic manifolds with some disjoint (open) horoballs removed. $\mathcal{H}_X$ and $\mathcal{H}_Y$ will denote the resulting horospheres.

Lemma 1.11. [32] *A Cannon-Thurston map for a weakly type-preserving proper embedding $i\colon Y \to X$ exists if and only if there exists a non-negative function $M(N)$ with $M(N) \to \infty$ as $N \to \infty$ such that the following holds:*
Suppose $y_0 \in Y$, and $\hat{\lambda}$ in $\widehat{Y} = \mathcal{E}(Y, \mathcal{H}_Y)$ is an electric quasigeodesic segment starting and ending outside horospheres. If $\lambda^b = \hat{\lambda} \setminus \bigcup_{K \in \mathcal{H}_Y} K$ lies outside an $B_N(y_0) \subset Y$, then for any electric quasigeodesic $\hat{\beta}$ joining the end points of $\hat{i}(\hat{\lambda})$ in $\widehat{X} = \mathcal{E}(X, \mathcal{H}_X)$, $\beta^b = \hat{\beta} \setminus \bigcup_{H \in \mathcal{H}_X} H$ lies outside $B_{M(N)}(i(y_0)) \subset X$.

We shall describe this informally as follows:
If λ lies outside a large ball **modulo horoballs** *then so does any geodesic in X joining its endpoints.*

In [11] we proved the existence of Cannon-Thurston maps for Kleinian groups corresponding to pared manifolds whose boundary is incompressible away from cusps.

Definition 1.12. *A* **pared manifold** *is a pair (M, P), where $P \subset \delta M$ is a (possibly empty) 2-dimensional submanifold with boundary such that*

1. *the fundamental group of each component of P injects into the fundamental group of M*
2. *the fundamental group of each component of P contains an abelian subgroup of finite index.*
3. *any cylinder $C : (S^1 \times I, \delta S^1 \times I) \to (M, P)$ such that $\pi_1(C)$ is injective is homotopic rel boundary into P.*
4. *P contains every component of δM which has an abelian subgroup of finite index.*

A pared manifold (M, P) is said to have **incompressible boundary** if each component of $\partial_0 M = \partial M \setminus P$ is incompressible in M.

The following Theorem summarizes the main results of [28], [29], [10], [11]. It proves the existence of Cannon-Thurston maps for Kleinian groups corresponding to pared manifolds whose boundary is incompressible away from cusps. It also describes the structure of these maps in terms of ending laminations.

Theorem 1.13. [11] *Suppose that $N^h \in H(M, P)$ is a hyperbolic structure on a pared manifold (M, P) with incompressible boundary. Let M_{gf} denotes a geometrically finite hyperbolic structure adapted to (M, P). Then the map $i : \widetilde{M_{gf}} \to \widetilde{N^h}$ extends continuously to the boundary $\partial i : \partial \widetilde{M_{gf}} \to \partial \widetilde{N^h}$.*

Let E be a degenerate end of N^h and $\widetilde{E}$ a lift of E to $\widetilde{N^h}$. Then the ending lamination $\mathcal{L}_E$ for the end E lifts to a lamination on $\widetilde{M_{gf}} \cap \widetilde{E}$. Each such lift $\mathcal{L}$ of the ending lamination of a degenerate end defines a relation $\mathcal{R}_{\mathcal{L}}$ on the (Gromov) hyperbolic boundary $\partial \widetilde{M_{gf}}$ given by $a \mathcal{R}_{\mathcal{L}} b$ iff a, b are end-points of a leaf of $\mathcal{L}$. Let $\{\mathcal{R}_i\}_i$ be the entire collection of relations on $\partial \widetilde{M_{gf}}$ obtained this way. Let $\mathcal{R}$ be the transitive closure of the union $\bigcup_i \mathcal{R}_i$. Then $\partial i(a) = \partial i(b)$ iff $a \mathcal{R} b$.

2. Split Geometry

Split level Surfaces

A **pants decomposition** of a compact surface S, possibly with boundary, is a disjoint collection of 3-holed spheres $P_1, \cdots, P_n$ embedded in S such that $S \setminus \bigcup_i P_i$ is a disjoint collection of non-peripheral annuli in S, no two of which are homotopic.

Let N be the convex core of a hyperbolic 3-manifold minus an open neighborhood of the cusp(s). Then any end E of N is simply degenerate [1], [14], [7] and homeomorphic to $S \times [0, \infty)$, where S is a compact surface, possibly with boundary. A closed geodesic in an end E homeomorphic to $S \times [0, \infty)$ is **unknotted** if it is isotopic in E to a simple closed curve in $S \times \{0\}$ via the homeomorphism. A **tube** in an end $E \subset N$ is a regular $R-$neighborhood $N(\gamma, R)$ of an unknotted geodesic γ in E.

Let $\mathcal{T}$ denote a collection of disjoint, uniformly separated tubes in ends of N such that

a) all Margulis tubes in E belong to $\mathcal{T}$ for all ends E of N.

b) there exists $\epsilon_0 > 0$ such that the injectivity radius $injrad_x(E) > \epsilon_0$ for all $x \in E \setminus \bigcup_{T\in\mathcal{T}} Int(T)$ and all ends E of N.

Let $F : N \to M$ be a bi-Lipschitz homeomorphism and let $M(0)$ be the image of $N \setminus \bigcup_{T\in\mathcal{T}} Int(T)$ in M under the bi-Lipschitz homeomorphism F. Let $\partial M(0)$ (resp. ∂M) denote the boundary of $M(0)$ (resp. M). M will be called the model manifold. The metrics on M and $\widetilde{M}$ will be denoted by d_M.

Let $(Q, \partial Q)$ be the unique hyperbolic pair of pants such that each component of ∂Q has length one. Q will be called the *standard* pair of pants. An isometrically embedded copy of $(Q, \partial Q)$ in $(M(0), \partial M(0))$ will be said to be *flat*.

Definition 2.1. A **split level surface** associated to a pants decomposition $\{Q_1, \cdots, Q_n\}$ of S in $M(0) \subset M$ is an embedding $f : \cup_i(Q_i, \partial Q_i) \to (M(0), \partial M(0))$ such that
1) Each $f(Q_i, \partial Q_i)$ is flat
2) f extends to an embedding (also denoted f) of S into M such that the interior of each annulus component of $f(S \setminus \bigcup_i Q_i)$ lies entirely in $F(\bigcup_{T\in\mathcal{T}} Int(T))$.

Let S_i^s denote the union of the collection of flat pairs of pants in the image of the embedding S_i.

The class of *all* topological embeddings from S to M that agree with a split level surface f associated to a pants decomposition $\{Q_1, \cdots, Q_n\}$ on $Q_1 \cup \cdots \cup Q_n$ will be denoted by $[f]$.

We define a partial order $\leq_E$ on the collection of split level surfaces in an end E of M as follows:
$f_1 \leq_E f_2$ if there exist $g_i \in [f_i]$, $i = 1, 2$, such that $g_2(S)$ lies in the unbounded component of $E \setminus g_1(S)$.

A sequence S_i of split level surfaces is said to exit an end E if $i < j$ implies $S_i \leq_E S_j$ and further for all compact subsets $B \subset E$, there exists $L > 0$ such that $S_i \cap B = \emptyset$ for all $i \geq L$.

Definition 2.2. *A curve v in $S \subset E$ is* ***l-thin*** *if the core curve of the Margulis tube $T_v(\subset E \subset N)$ has length less than or equal to l. A tube $T \in \mathcal{T}$ is l-thin if its core curve is l-thin. A tube $T \in \mathcal{T}$ is l-thick if it is not l-thin.*
A curve v is said to split a pair of split level surfaces S_i and S_j $(i < j)$ if v occurs as a boundary curve of both S_i and S_j.

The collection of all l-thin tubes is denoted as $\mathcal{T}_l$. The union of all l-thick tubes with $M(0)$ is denoted as $M(l)$.

Definition 2.3. *A pair of split level surfaces S_i and S_j $(i < j)$ is said to be* ***k-separated*** *if*
a) for all $x \in S_i^s$, $d_M(x, S_j^s) \geq k$
b)Similarly, for all $x \in S_j^s$, $d_M(x, S_i^s) \geq k$.

Definition 2.4. An L-bi-Lipschitz **split surface** in $M(l)$ associated to a pants decomposition $\{Q_1, \cdots, Q_n\}$ of S and a collection $\{A_1, \cdots, A_m\}$ of complementary annuli in S is an embedding $f : \cup_i Q_i \bigcup \cup_i A_i \to M(l)$ such that
1) the restriction $f : \cup_i(Q_i, \partial Q_i) \to (M(0), \partial M(0))$ is a split level surface
2) the restriction $f : A_i \to M(l)$ is an L-bi-Lipschitz embedding.

3) f extends to an embedding (also denoted f) of S into M such that the interior of each annulus component of $f(S \setminus (\cup_i Q_i \bigcup \cup_i A_i))$ lies entirely in $F(\bigcup_{T \in \mathcal{T}_l} Int(T))$.

Note: The difference between a split level surface and a split surface is that the latter may contain bi-Lipschitz annuli in addition to flat pairs of pants.

We denote split surfaces by Σ_i to distinguish them from split level surfaces S_i. Let Σ_i^s denote the union of the collection of flat pairs of pants and bi-Lipschitz annuli in the image of the split surface (embedding) Σ_i.

Theorem 2.5. *Let $N, M, M(0), S, F$ be as above and E an end of M. For any l less than the Margulis constant, let $M(l) = \{F(x) : \mathrm{injrad_x}(N) \geq l\}$. Fix a hyperbolic metric on S such that each component of ∂S is totally geodesic of length one (this is a normalization condition). There exist $L_1 \geq 1$, $\epsilon_1 > 0$, $n \in \mathbb{N}$, and a sequence Σ_i of L_1-bi-Lipschitz, ϵ_1-separated split surfaces exiting the end E of M such that for all i, one of the following occurs:*

1. *An l-thin curve v splits the pair (Σ_i, Σ_{i+1}), i.e. v splits the associated split level surfaces (S_i, S_{i+1}), which in turn form an l-thin pair.*
2. *there exists an L_1-bi-Lipschitz embedding*
$$G_i : (S \times [0,1], (\partial S) \times [0,1]) \to (M, \partial M)$$
such that $\Sigma_i^s = G_i(S \times \{0\})$ *and* $\Sigma_{i+1}^s = G_i(S \times \{1\})$

Finally, each l-thin curve in S splits at most n split level surfaces in the sequence $\{\Sigma_i\}$.

A model manifold M all of whose ends are equipped with a collection of exiting split surfaces satisfying the conclusions of Theorem 2.5 is said to be equipped with a **weak split geometry** structure.

Pairs of split surfaces satisfying Alternative (1) of Theorem 2.5 will be called an l-thin pair of split surfaces (or simply a thin pair if l is understood). Similarly, pairs of split surfaces satisfying Alternative (2) of Theorem 2.5 will be called an l-thick pair (or simply a thick pair) of split surfaces.

Definition 2.6. *Let $(\Sigma_i^s, \Sigma_{i+1}^s)$ be a thick pair of split surfaces in M. The closure of the bounded component of $M \setminus (\Sigma_i^s \cup \Sigma_{i+1}^s)$ between $\Sigma_i^s, \Sigma_{i+1}^s$ will be called a thick block.*

Note that a thick block is uniformly bi-Lipschitz to the product $S \times [0,1]$ and that its boundary components are $\Sigma_i^s, \Sigma_{i+1}^s$.

Definition 2.7. *Let $(\Sigma_i^s, \Sigma_{i+1}^s)$ be an l-thin pair of split surfaces in M and $F(\mathcal{T}_i)$ be the collection of l-thin Margulis tubes that split both $\Sigma_i^s, \Sigma_{i+1}^s$. The closure of the union of the bounded components of $M \setminus ((\Sigma_i^s \cup \Sigma_{i+1}^s) \bigcup_{F(T) \in F(\mathcal{T}_i)} F(T))$ between $\Sigma_i^s, \Sigma_{i+1}^s$ will be called a split block. Equivalently, the closure of the union of the bounded components of $M(l) \setminus (\Sigma_i^s \cup \Sigma_{i+1}^s)$ between $\Sigma_i^s, \Sigma_{i+1}^s$ is a split block. Each connected component a split block is a split component.*

Remark 2.8. For each lift $\widetilde{K} \subset \widetilde{M}$ of a split component K of a split block of $M(l) \subset M$, there are lifts of l-thin Margulis tubes that share the boundary of $\widetilde{K}$ in $\widetilde{M}$. Adjoining these lifts to $\widetilde{K}$ we obtain **extended split components**. Let $\mathcal{K}'$ denote the collection of extended split components in $\widetilde{M}$. Denote the collection of split components in $\widetilde{M(l)} \subset \widetilde{M}$ by $\mathcal{K}$. Let $\widetilde{M(l)}$ denote the lift of $M(l)$ to $\widetilde{M}$. Then the inclusion of $\widetilde{M(l)}$ into $\widetilde{M}$ gives a quasi-isometry between $\mathcal{E}(\widetilde{M(l)}, \mathcal{K})$ and $\mathcal{E}(\widetilde{M}, \mathcal{K}')$ equipped with the respective electric metrics. This follows from the last assertion of Theorem 2.5.

The electric metric on $\mathcal{E}(\widetilde{M}, \mathcal{K}')$ is called the **graph-metric** and is denoted by d_G. The electric space will be denoted as $(\widetilde{M}, d_G)$.

The electric metric on $\mathcal{E}(\widetilde{M}, \mathcal{K} \bigcup \mathcal{T}_l)$ is quasi-isometric to the electric metric on $\mathcal{E}(\widetilde{M}, \mathcal{K}')$, again by the last assertion of Theorem 2.5. The electric space will be denoted as $(\widetilde{M}, d_G^1)$.

Definition 2.9. *Let $Y \subset \widetilde{N}$ and $X = F(Y)$. $X \subset \widetilde{M}$ is said to be Δ-graph quasiconvex if for any hyperbolic geodesic μ joining $a, b \in Y$, $F(\mu)$ lies inside $N_\Delta(X, d_G) \subset \mathcal{E}(\widetilde{M}, \mathcal{K}')$.*

For X a split component in a manifold, define $CH(X) = F(CH(Y))$, where $CH(Y)$ is the convex hull of Y in $\widetilde{N}$, provided the ends of N have no cusps, i.e. $N = N^h$. Else define $CH(X)$ to be the image under F of $CH(Y)$ minus cusps. Further, in order to ensure hyperbolicity of the universal cover, we partially electrocute the cusps of M (cf. Theorem 1.8).

Then Δ-graph quasiconvexity of X is equivalent to the condition that $dia_G(CH(X))$ is bounded by $\Delta' = \Delta'(\Delta)$ as any split component has diameter one in $(\widetilde{M}, d_G)$.

A split component $K (\subset E) \subset N$ is incompressible if the map $i_* : \pi_1(K) \to \pi_1(N)$ induced by the inclusion is injective. Lemma 2.10, Proposition 2.11 and Proposition 2.12 below were proved in [28] for M homotopy equivalent to a surface, where all split components are automatically incompressible. However the proofs in [28] require only that the split components be incompressible in M.

Lemma 2.10. *Let E be a simply degenerate end of a hyperbolic 3-manifold N equipped with a weak split geometry model M. For K an incompressible split component contained in E, let $\tilde{K}$ be a lift to $\tilde{N}$. Then there exists $C_0 = C_0(K)$ such that the convex hull of $\tilde{K}$ minus cusps lies in a C_0-neighborhood of $\tilde{K}$ in $\tilde{N}$.*

Proposition 2.11. *If K is an incompressible split component, then $\widetilde{K}$ is uniformly graph-quasiconvex in $\widetilde{M}$, i.e. there exists Δ' such that $dia_G(CH(\widetilde{K}))) \leq \Delta^prime$ for all incompressible split components $\widetilde{K}$.*

Proposition 2.12. *Suppose that all split components of $\widetilde{M}$ are incompressible. Then $(\widetilde{M}, d_G)$ and hence $(\widetilde{M}, d_G^1)$ are Gromov-hyperbolic.*

In fact electro-ambient quasigeodesics in $(\widetilde{M}, d_G)$ and $(\widetilde{M}, d_G^1)$ have the following relation.

Lemma 2.13. *Let M be the a model of split geometry such that all split components are incompressible. Let $(\widetilde{M}, d_G)(= \mathcal{E}(\widetilde{M}, \mathcal{K}'))$ and $(\widetilde{M}, d_G^1)(= \mathcal{E}(\widetilde{M}, \mathcal{K} \bigcup \mathcal{T}_l))$ be as above. Given $o \in \widetilde{M}$ and $C_0 > 0$, there exists a function $\Theta : \mathbb{N} \to \mathbb{N}$ satisfying $\Theta(n) \to \infty$ as $n \to \infty$ such that the following holds.*
For any $a, b \in \widetilde{M}$, let β_{ea}^h be an electro-ambient C_0-quasigeodesic without backtracking in $(\widetilde{M}, d_G)$ joining a, b. Let $\beta_{ea} = \beta_{ea}^h \setminus \partial\widetilde{M}$ be the part of β_{ea}^h lying away from the (bi-Lipschitz) horospherical boundary of $\widetilde{M}$. Again, let β_{ea1}^h be an

electro-ambient C_0-*quasigeodesic without backtracking in* $(\widetilde{M}, d_G^1)$ *joining* a, b. *Let* $\beta_{ea1} = \beta_{ea1}^h \setminus \partial\widetilde{M}$ *be the part of* β_{ea1}^h *lying away from the (bi-Lipschitz) horospherical boundary of* $\widetilde{M}$.

Then $d_M(\beta_{ea}, o) \geq n$ *implies that* $d_M(\beta_{ea1}, o) \geq \Theta(n)$. *Conversely,* $d_M(\beta_{ea1}, o) \geq n$ *implies that* $d_M(\beta_{ea}, o) \geq \Theta(n)$.

Proof. Let K' be an extended split component in $\mathcal{K}'$ and $\widetilde{K'}$ denote its universal cover. Let $\widetilde{K'} = \widetilde{K} \bigcup \widetilde{T^i}$ where $\widetilde{T^i}$ are the universal covers of l-thin Margulis tubes abutting the split component $\widetilde{K}$ in $\widetilde{K'}$. Suppose $K \subset B_i$ the i-th split block in an end E.

Then $\widetilde{K'}$ is hyperbolic and contained in a $C(i)$-neighborhood of $\widetilde{K}$. The argument is now a reprise of similar arguments in Section 6 of [28].

Hence for all i, there exists $C(i)$, such that $\beta_{ea1} \cap \widetilde{B_i}$ lies in a $C(i)$-neighborhood of $\beta_{ea} \cap \widetilde{B_i}$ in $\widetilde{M}$. Suppose $d_M(\beta_{ea}, o) \geq n$. Hence, by uniform k_0-separatedness of split surfaces, $d_M(\beta_{ea1} \cap \widetilde{B_i}, o) \geq max(n - C(i), ik_0)$.

Let $D(i) = max_{1 \leq j \leq i} C(i)$. Then $d_M(\beta_{ea1}, o) \geq max(n - D(i), ik_0)$ for all i. The Lemma follows.

The converse direction is similar. $\square$

We summarize the conclusions of the above propositions below.

Definition 2.14. *A model manifold of weak split geometry is said to be of* **split geometry** *if*

1. *Each split component* $\widetilde{K}$ *is quasiconvex (not necessarily uniformly) in the hyperbolic metric on* $\widetilde{N}$.
2. *Equip* $\widetilde{M}$ *with the graph-metric* d_G *obtained by electrocuting (extended) split components* $\widetilde{K}$. *Then the convex hull* $CH(\widetilde{K})$ *of any split component* $\widetilde{K}$ *has uniformly bounded diameter in the metric* d_G.

Hence by Lemma 2.10 and Proposition 2.11 we have the following.

Theorem 2.15. *Any degenerate end of a hyperbolic 3-manifold is bi-Lipschitz homeomorphic to a Minsky model and hence to a model of split geometry.*

3. Free Groups and Finitely Generated Kleinian Groups

Let G be a free geometrically infinite Kleinian group. Agol [1], and independently, Gabai and Calegari [14] have shown that $N = \mathbf{H}^3/G$ is topologically tame, and hence, by work of Canary [7], geometrically tame. Then any manifold M bi-Lipschitz to N is homeomorphic to the interior of a handlebody with boundary S.

More generally, let G be a finitely generated Kleinian group and G_f be a geometrically finite Kleinian group, abstractly isomorphic to G via a type-preserving isomorphism. Let H denote the convex core of $\mathbf{H}^3/G_f$ and let M be bi-Lipschitz homeomorphic to $N = \mathbf{H}^3/G$. Then there is a natural identification $i : H \to M$ of H with the augmented Scott core (i.e. Scott core plus parabolics) of M. Let $\widetilde{i}$ indicate the lift of i to the universal cover. For most of the discussion below, it might be helpful at a first reading to have in mind a free geometrically infinite

Kleinian group without parabolics. We fix this notation for H, M, S throughout this section.

3.1. **The Masur Domain.**

Definition 3.1. *Let E be an end of a hyperbolic manifold such that E is homeomorphic to $S \times [0, \infty)$ for S a finite area hyperbolic surface. A map $h : E \to S \times [0, \infty)$ is said to be type-preserving, if all and only the cusps of E are mapped to cusps of $S \times [0, \infty)$.*

Theorem 3.2. [1] [14] [3] [5] *Let G be a finitely generated Kleinian group and $M = \mathbf{H}^3/G$. Let H denote an augmented Scott core of M. Let E_1 be a geometrically infinite end of $M \setminus H$. Then E_1 is homeomorphic (via a type-preserving homeomorphism) to a topological product $S \times [0, \infty)$ for a hyperbolic surface S of finite area. Further, there exists a neighborhood E of the end corresponding to E_1 such that E is bi-Lipschitz homeomorphic to a Minsky model for $S \times [0, \infty)$ and hence to a model of split geometry.*

The last part of the last statement follows from Theorem 2.15.

Some ambiguity remains in the statement of Theorem 3.2 above. This lies in the choice of the ending lamination for E used to build the Minsky model. Since $i : S \subset E$ is type-preserving, no parabolic element of S bounds a compressing disk.

Let $\mathcal{ML}(S)$ be the space of measured laminations. Let $\mathcal{D}(S)$ be the subset of $\mathcal{ML}(S)$ consisting of weighted unions of disjoint meridians (i.e. boundaries of compression disks lying on S). Let $cl(\mathcal{D}(S))$ denote the closure of $\mathcal{D}(S)$. Define the **Masur domain** of S by

$\mathcal{MD}(S) = \{\lambda \in \mathcal{ML}(S) : i(\lambda, \mu) > 0\}$ for all $\mu \in cl(\mathcal{D}(S))$, provided S has at least two disjoint isotopy classes of compressing disks.

Else, we define

$\mathcal{MD}(S) = \{\lambda \in \mathcal{ML}(S) : i(\lambda, \mu) > 0$ for any μ that is disjoint from a compressing disk $\}$

Now, let $M = H \cup_i E_i$, where H is an augmented Scott core. Let $S = H \cap E$ be one of the boundary components of H. Let $Mod_0(S)$ denote the subgroup of the mapping class group of S generated by Dehn twists along essential simple closed curves that bound embedded disks in H. As is customary, a prefix $\mathcal{P}$ will indicate projectivization. $Mod_0(S)$ acts on $\mathcal{PML}(S)$. It was shown by Otal [33] (see also McCarthy and Papadopoulos [20]) that under this action, $Mod_0(S)$ acts properly discontinuously on $\mathcal{PMD}(S)(\subset \mathcal{PML}(S))$ with limit set $Pcl(\mathcal{D}(S))(\subset \mathcal{PML}(S))$.

The ending lamination is well-defined up to the action of $Mod_0(S)$ (see [7]).

Theorem 3.3. [7] *For any finitely generated Kleinian group, the ending lamination λ facing a surface S with a compressing disk lies in the Masur Domain.*

For our purposes we shall mostly be satisfied with the fact that E is bi-Lipschitz homeomorphic to *some* Minsky model, and hence, by Theorem 3.2.

3.2. **Incompressibility of Split Components.**

We would like to show that sufficiently deep within an end E, all split components are incompressible *in* M. Recall that splitting tubes correspond to *thin Margulis tubes* in the split geometry model built from the Minsky model.

Proposition 3.4. *Let $M = H \cup E$. Equip E with a split geometry structure. Then there exists $E_2 \subset E$ such that*

1. *E_2 is homeomorphic to $S\times[0,\infty)$ by a type-preserving homeomorphism and consists of a union of blocks and tubes from the split geometry model for E.*
2. *All split components of E_2 are incompressible, i.e. if K is a split component of E_2, then the inclusion $i : K \to M$ induces an injective map $i_* : \pi_1(K) \to \pi_1(M)$.*

Proof: Suppose not. Then there exists a sequence of split components K_i exiting the end E such that $i_* : \pi_1(K) \to \pi_1(M)$ is not injective. Since K_i are split components, $K_i = S_i \times I$ for some subsurface S_i of S. By the Loop Theorem (see for instance, Hempel [15]), there exist simple closed curves $\sigma_i \subset S_i$ such that σ_i bound embedded topological disks in H. Hence $\sigma_i \in \mathcal{D}(S)$. Let T_i be a splitting tube bounding K_i. Then T_i has a core curve α_i, which in turn corresponds to a simple closed curve on S. It follows that $d_{CC}(\alpha_i, \mathcal{D}(S)) = 1$, where d_{CC} denotes distance in the curve-complex. Since any such sequence of curves α_i converges to the ending lamination λ corresponding to the end E, it follows that $\lambda \in cl(\mathcal{D}(S))$ and hence cannot lie in the Masur domain. This contradicts Theorem 3.3. The proposition follows. □

As an immediate consequence we have the following.

Lemma 3.5. *If K is a split component, then $\pi_1(K)(\subset \pi_1(M))$ is geometrically finite (Schottky, in the absence of parabolics).*

Proof. Follows from Lemma 2.10. □

Proposition 3.6. *If K is a split component, then $\widetilde{K}$ is uniformly graph-quasiconvex in $\widetilde{M}$.*

Proof. Follows from Proposition 2.11. □

Proposition 3.7. *$(\widetilde{M}, d_G)$ is a hyperbolic metric space.*

Proof. Follows from Proposition 2.12. □

3.3. **Constructing Quasidisks.**

The construction in this subsection may be regarded as a *graph-metrized coarse* analogue of an unpublished construction due to Miyachi [27] (see also Souto [34]). The main technical difference between Miyachi's construction and ours is that Miyachi constructs continuous images of disks that actually separate the universal cover $\widetilde{M}$, whereas we only construct quasidisks. As a consequence it becomes technically more difficult for us to prove that quasidisks coarsely separate. This is why we need a special family of paths which we shall call 'admissible paths' in the next subsection which either intersect or come close to the quasidisks we construct below.

We choose a collection of essential simple closed curves $\sigma_1 \cdots \sigma_g$ on S bounding disks $D_1 \cdots D_g$ with neighborhoods $D_i \times (-\epsilon, \epsilon)$ such that each component of $H \setminus \bigcup_i D_i \times (-\epsilon, \epsilon)$ is either a ball or has incompressible boundary (rel. cusps). Also assume that σ_i are geodesics in the intrinsic metric on S. Next, let E be described as a union of contiguous blocks B_k, where each B_k is either a split block, or a thick block.

Further, let $\partial B_k = S_{k-1} \cup S_k$ with S_k the *upper boundary* and S_{k-1} the *lower boundary.* Also let $S = S_0$ and $\sigma_i = \sigma_{i0}$. Let σ_{ik} be the shortest closed curve in the split metric on S_k (i.e. in the pseudometric obtained by electrocuting annular intersections of splitting Margulis tubes with the split level surface S_k) homotopic in E to $\sigma_i = \sigma_{i0}$. Let $\overline{A_i} = D_i \bigcup_k \sigma_{ik} \subset M$ be the union of the disk D_i and the quasi-annulus $\bigcup_k \sigma_{ik}$. Then any lift A_i of $\overline{A_i}$ to $\widetilde{M}$ is isometric to $\overline{A_i}$ as D_i is homotopically trivial and σ_{ik} are all freely homotopic to $\sigma_i = \sigma_{i0} = \partial D_i$.

We want to show that A_i are quasiconvex in $(\widetilde{M}, d_G)$ which is hyperbolic by Proposition 3.7.

qi Rays

Fix a σ and the disk D it bounds. Let $A = D \bigcup_k \sigma_k \subset M$, where $\sigma_k \subset S_k$. Lift $\bigcup_k \sigma_k$ to the universal cover $\widetilde{\widetilde{E}}$ of E such that any lift $\widetilde{\sigma_k}$ lies in the universal cover $\widetilde{\widetilde{S_k}}$ of S_k (We are using this notation to distinguish from lifts to $\widetilde{M}$). Let λ_k be any such lift $\widetilde{\sigma_k}$. We then have the following from [28].

Lemma 3.8. [28] *There exists $C \geq 0$ such that for $x_k \in \lambda_k$ there exists $x_{k-1} \in \lambda_{k-1}$ with $d_G(x_k, x_{k-1}) \leq C$. Similarly there exists $x_{k+1} \in \lambda_{k+1}$ with $d_G(x_k, x_{k+1}) \leq C$. Also, for all k there exists $B(k)$ such that for all $x_k \in \lambda_k$ there exists $x_{k-1} \in \lambda_{k-1}$ with $d_M(x_k, x_{k-1}) \leq B(k)$. Hence, for all n and $x \in \lambda_n$, there exists a C-quasigeodesic ray r such that $r(k) \in \lambda_k \subset \mathcal{L}_\lambda$ for all k and $r(n) = x$.*

Further, by construction of split blocks, $d_G(x_i, S_{i-1}) = 1$. Therefore inductively, $d_G(x_i, S_j) = |i-j|$. Hence $d_G(x_i, x_j) \geq |i-j|$. By construction, $d_G(x_i, x_j) \leq C|i-j|$.

Hence, given $p \in \lambda_i$ the sequence of points $x_n, n \in \mathbb{N} \cup \{0\}$ with $x_i = p$ gives by Lemma 3.8 above, a quasigeodesic in the d_G-metric. Such quasigeodesics shall be referred to as *d_G-quasigeodesic rays.*

After projecting $\widetilde{E}$ to $\widetilde{M} \setminus \widetilde{H}$ we have the following conclusion.

Lemma 3.9. *There exists $C \geq 0$ such that for all k there exists B_k satisfying the following:*
For all $x_{ik} \in \sigma_{ik}$ there exists $x_{i,k-1} \in \sigma_{i,k-1}$ with $d_G(x_{ik}, x_{i,k-1}) \leq C$ and $d(x_{ik}, x_{i,k-1}) \leq B_k$.

The following Corollary will turn out to be quite useful.

Corollary 3.10. *There exists $C \geq 0$ such that for all k and all $x_{ik} \in \sigma_{ik}$ there exists $q \in \sigma_{i0}$ and a sequence of points $p = x_{ik}, \cdots, x_{i0} = q$ which is a quasigeodesic in $(\widetilde{M}, d_G)$.*

Proof. By construction of split blocks, $d_G(x_{ik}, S_{k-1}) = 1$.

Hence, given $p \in \sigma_{ik}$ the sequence of points $p = x_{ik}, \cdots, x_{i0}$ gives by Lemma 3.9 above, a quasigeodesic in the d_G-metric lying entirely on A_i joining p to a point $q \in D_i$. □

We can choose a point $z_i \in D_i$ (quite arbitrarily) and extend any quasigeodesic constructed as above by adding on a path from q to z_i lying entirely in D_i and having uniformly bounded length (This can be done easily as D_i has bounded diameter.)

Proposition 3.11. *There exists $C_0 \geq 0$ such that each A_j is C_0-quasiconvex in $(\widetilde{M}, d_G)$.*

Proof: By Lemma 3.9 and Corollary 3.10 above, it follows that there exist $K \geq 1$ such that for any two points p_1, p_2 in A_j there exist K- quasi-geodesics γ_1, γ_2 to z_j.

By Proposition 3.7 we also have that $(\widetilde{M}, d_G)$ is hyperbolic. Hence any geodesic α_i $(i = 1, 2)$ joining p_i to z_j lies in some K_1 neighborhood of γ_i. Further, by hyperbolicity of $(\widetilde{M}, d_G)$, we conclude that a geodesic β joining p_1, p_2 lies in a K_2-neighborhood of $\alpha_1 \cup \alpha_2$. Hence, finally, β lies in a $(K_1 + K_2)$ neighborhood of $\gamma_1 \cup \gamma_2 \subset A_j$. Choosing $C_0 = K_1 + K_2$, we are through. □

The quasidisks constructed above have the following property.

Lemma 3.12. *Let M be a model manifold of split geometry. Let Q be a Scott core of M and $\{D_i\}$ a maximal collection of compressing disks in Q. Then there exists a function $\Theta : \mathbb{N} \to \mathbb{N}$ satisfying $\Theta(n) \to \infty$ as $n \to \infty$ such that for all $o \in \widetilde{Q}$ the following holds.*
Let D be a lift of one of the D_i's to $\widetilde{M}$ and let A be the quasidisk in $\widetilde{M}$ constructed from D as above.

Then $d_M(D, o) \geq n$ implies that $d_M(A, o) \geq \Theta(n)$.

Proof. By Lemma 3.9 and Corollary 3.10, there exists $b_1, \cdots, b_k, \cdots$ and $z \in D$ such that for all $x_k \in \sigma_k \subset A$, $d_M(x_k, z) \leq (b_1 + \cdots + b_k) = c_k$(say). Hence $d_M(x_k, 0) \geq (n - c_k)$.

By uniform ϵ_0-separatedness of split surfaces, $d_M(x_k, 0) \geq k\epsilon_0$. Hence $d_M(x_k, 0) \geq max((n - c_k), k\epsilon_0)$. Choosing $\Theta(n)$ to be the largest value of k such that $k\epsilon_0 \leq n - c_k$ we are done. □

3.4. **Reduction Lemma and Admissible Paths.**

Before we get into the proof of the existence of Cannon-Thurston maps, we recall some material from Section 6 of [28] that will help streamline the proof.

The next Lemma allows us to apply the criterion for existence of Cannon-Thurston maps in Lemmas 1.10and 1.11 to electro-ambient quasigeodesics in $\widetilde{M}$ rather than hyperbolic geodesics in $\widetilde{N}$. Lemma 3.13 below is a paraphrasing of what Lemmas 6.8 and 6.9 of [28] prove.

Lemma 3.13. [28] *Let N be the convex core of a complete hyperbolic $3-$manifold N^h minus a neighborhood of the cusps. Equip each degenerate end with a split geometry structure such that each split component is incompressible. Let M be the resulting model of split geometry and $F : N \to M$ be the bi-Lipschitz homeomorphism between the two. Let $\widetilde{F}$ be a lift of F to the universal covers. Then for all $C_0 > 0$, and $o \in \widetilde{N}$ there exists a function $\Theta : \mathbb{N} \to \mathbb{N}$ satisfying $\Theta(n) \to \infty$ as $n \to \infty$ such that the following holds.*
For any $a, b \in \widetilde{N} \subset \widetilde{N^h}$, let λ^h be the hyperbolic geodesic in $\widetilde{N^h}$ joining them and let $\lambda^h_{thick} = \lambda^h \cap \widetilde{N}$. Similarly let β^h_{ea} be an electro-ambient C_0-quasigeodesic without backtracking in $\widetilde{M} \subset \mathcal{E}(\widetilde{M}, \mathcal{K}')$ joining $\widetilde{F}(a)$, $\widetilde{F}(b)$. Let $\beta_{ea} = \beta^h_{ea} \setminus \partial\widetilde{M}$ be the part of β^h_{ea} lying away from the (bi-Lipschitz) horospherical boundary of $\widetilde{M}$.

Then $d_M(\beta_{ea}, \widetilde{F}(o)) \geq n$ implies that $d_{\mathbf{H}^3}(\lambda^h_{thick}, o) \geq \Theta(n)$.

Combining Lemma 3.13 with Lemma 2.13, we have the following.

Corollary 3.14. *Let N be the convex core of a complete hyperbolic $3-$manifold N^h minus a neighborhood of the cusps. Equip each degenerate end with a split geometry structure such that each split component is incompressible. Let M be the resulting model of split geometry and $F : N \to M$ be the bi-Lipschitz homeomorphism between the two. Let $\widetilde{F}$ be a lift of F to the universal covers. Then for all $C_0 > 0$, and*

$o \in \widetilde{N}$ there exists a function $\Theta : \mathbb{N} \to \mathbb{N}$ satisfying $\Theta(n) \to \infty$ as $n \to \infty$ such that the following holds.
For any $a, b \in \widetilde{N} \subset \widetilde{N^h}$, let λ^h be the hyperbolic geodesic in $\widetilde{N^h}$ joining them and let $\lambda^h_{thick} = \lambda^h \cap \widetilde{N}$. Similarly let β^h_{ea} be an electro-ambient C_0-quasigeodesic without backtracking in $\widetilde{M} \subset \mathcal{E}(\widetilde{M}, \mathcal{K} \bigcup \mathcal{T}_l)$ joining $\widetilde{F}(a)$, $\widetilde{F}(b)$. Let $\beta_{ea} = \beta^h_{ea} \setminus \partial\widetilde{M}$ be the part of β^h_{ea} lying away from the (bi-Lipschitz) horospherical boundary of $\widetilde{M}$.

Then $d_M(\beta_{ea}, \widetilde{F}(o)) \geq n$ implies that $d_{\mathbf{H}^3}(\lambda^h_{thick}, o) \geq \Theta(n)$.

Again, combining Lemma 3.13 with Lemma 3.12, we have the following.

Corollary 3.15. *Let N be the convex core of a complete hyperbolic $3-$manifold N^h minus a neighborhood of the cusps. Equip each degenerate end with a split geometry structure such that each split component is incompressible. Let M be the resulting model of split geometry and $F : N \to M$ be the bi-Lipschitz homeomorphism between the two. Let $\widetilde{F}$ be a lift of F to the universal covers. Let Q be a Scott core of N and $\{D_i\}$ a maximal collection of compressing disks in Q. Then there exists a function $\Theta : \mathbb{N} \to \mathbb{N}$ satisfying $\Theta(n) \to \infty$ as $n \to \infty$ such that for all $o \in \widetilde{Q}$ the following holds.*
Let D be a lift of one of the D_i's to $\widetilde{M}$ and let A be the quasidisk in $\widetilde{M}$ constructed from D as above. For any $a, b \in A$, let $[a, b]_h$ be the hyperbolic geodesic in $\widetilde{N^h}$ joining $\widetilde{F}^{-1}(a), \widetilde{F}^{-1}(b)$ and let $[a, b] = [a, b]_h \cap \widetilde{N}$.

Then $d_M(D, \widetilde{F}(o)) \geq n$ implies that $d_{\mathbf{H}^3}([a, b], o) \geq \Theta(n)$.

Proof. By Lemma 3.12 there exists $z \in D$, a function $\Theta_0 : \mathbb{N} \to \mathbb{N}$ satisfying $\Theta_0(n) \to \infty$ as $n \to \infty$ and electro-ambient quasigeodesics β_a, β_b in $(\widetilde{M}, d_G)$ such that $d_M(D, \widetilde{F}(o)) \geq n$ implies that $d_M(\beta_a \cup \beta_b, \widetilde{F}(o)) \geq \Theta_0(n)$.

Let $[a, z]_h, [b, z]_h$ be the hyperbolic geodesic in $\widetilde{N}$ joining $\widetilde{F}^{-1}(a), \widetilde{F}^{-1}(b)$ respectively to $\widetilde{F}^{-1}(z)$ and let $[a, z] = [a, z]_h \cap \widetilde{N}$, $[b, z] = [b, z]_h \cap \widetilde{N}$.

Then by Lemma 3.13 there exists a function $\Theta_1 : \mathbb{N} \to \mathbb{N}$ satisfying $\Theta_1(n) \to \infty$ as $n \to \infty$ such that $d_M(D, \widetilde{F}(o)) \geq n$ implies that $d_{\mathbf{H}^3}([a, z] \cup [b, z], o) \geq \Theta_1(n)$.

Let $\delta > 0$ be such that all geodesic triangles in $\mathbb{H}^3$ are $\delta-$thin. Taking $\Theta(n) = \Theta_1(n) - \delta$, it follows that $d_M(D, \widetilde{F}(o)) \geq n$ implies that $d_{\mathbf{H}^3}([a, b], o) \geq \Theta(n)$. □

Admissible Quasigeodesics
We shall need a collection of paths consisting of horizontal and vertical segments approximating electro-ambient quasigeodesics. We shall call these admissible quasigeodesics. Let M be a model manifold each of whose ends is equipped with a split geometry structure such that all split components are incompressible. Recall that each thick block and each split block in M is homeomorphic to a product $\Sigma^s_i \times I$. We fix such a product structure for each block. Let $t_i = sup\{length(\{x\} \times I) : x \in \Sigma^s_i\}$ be the thickness of the $i-$th block.

Recall that $F : N \to M$ is a bi-Lipschitz homeomorphism from a hyperbolic manifold N (minus cusps) to M and let $\widetilde{F}$ denote its lift to the universal cover.

An **elementary admissible path** in $\widetilde{M}$ is one of the following:

1. A 'horizontal' geodesic in the intrinsic path metric on some lift $\widetilde{\Sigma^s_i}$ of a split surface to $\widetilde{M}$.
2. A 'vertical' path of the form $x \times I$ (with respect to the fixed product structure above) in the lift to $\widetilde{M}$ of either a thick block or a split block.

Let B_i be a thick block of M and let S^s_i, S^s_{i+1} be its horizontal boundary components. Recall that a product structure $B_i = S \times I$ has been fixed. Let $\mu = [a, b]$ be a geodesic in $\widetilde{B_i} \subset \widetilde{M}$ such that its end-points a, b lie on the horizontal boundary components. Let $P(\mu)$ denote the projection of μ onto the horizontal boundary component (S^s_i or S^s_{i+1}) containing a. If b belongs to the same horizontal boundary component as a, define $\mu_{adm} = P(\mu)$. Else define $\mu_{adm} = P(\mu) \cup \{b\} \times I$, where $\{b\} \times I$ is the elementary vertical path through b. μ_{adm} will be called the **admissible quasigeodesic** corresponding to μ in the thick block $\widetilde{B_i}$.

Let $\mathcal{B}$ denote the collection of thick blocks.

Definition 3.16. *An* **admissible quasigeodesic** β_{adm} *in* $\widetilde{M}$ *corresponding to an electro-ambient* C_0-*quasigeodesic* β_{ea} *without backtracking in* $\widetilde{M} \subset \mathcal{E}(\widetilde{M}, \mathcal{K} \bigcup \mathcal{T}_l)$ *is a path such that* $\beta_{adm} \cap (\widetilde{M} \setminus \bigcup_{K \in \mathcal{K}} \widetilde{K} \cup \bigcup_{B \in \mathcal{B}} \widetilde{B}) = \beta_{ea} \cap (\widetilde{M} \setminus \bigcup_{K \in \mathcal{K}} \widetilde{K} \cup \bigcup_{B \in \mathcal{B}} \widetilde{B})$.

Further, for each $\widetilde{K}$, $\beta_{adm} \cap \widetilde{K}$ *is a union of elementary admissible paths with disjoint interiors such that*

1. $\beta_{adm} \cap \widetilde{K}$ *has at most one vertical path of type (2) above.*
2. *for any connected horizontal boundary component* $\widetilde{\Sigma^s_0}$ *of* $\widetilde{K}$, $\beta \cap \widetilde{\Sigma^s_0}$ *has at most one 'horizontal' geodesic of type (1) above.*

Finally for each $B \in \mathcal{B}$, $\beta_{adm} \cap \widetilde{B}$ *is the admissible quasigeodesic corresponding to* $\beta_{ea} \cap \widetilde{B}$ *in* $\widetilde{B}$.

The next Lemma allows us to apply the criterion for existence of Cannon-Thurston maps in Lemma 1.10 to admissible quasigeodesics in $\widetilde{M}$ rather than electro-ambient quasigeodesics in $\widetilde{M}$. The proof of Lemma 3.17 is exactly like Lemma 6.5 of [28] and we omit it here (see also the proof of Lemma 2.13 above).

Lemma 3.17. [28] *Let* N *be the convex core of a complete hyperbolic* $3-$*manifold* N^h *minus a neighborhood of the cusps. Equip each degenerate end with a split geometry structure such that each split component is incompressible. Let* M *be the resulting model of split geometry. Then for all* $C_0 > 0$, *and* $o \in \widetilde{M}$ *there exists a function* $\Theta : \mathbb{N} \to \mathbb{N}$ *satisfying* $\Theta(n) \to \infty$ *as* $n \to \infty$ *such that the following holds. For any* $a, b \in \widetilde{M}$, *let* β^h_{ea} *be an electro-ambient* C_0-*quasigeodesic without backtracking in* $\widetilde{M}$ *joining* a, b. *Let* $\beta_{ea} = \beta^h_{ea} \setminus \partial\widetilde{M}$ *be the part of* β^h_{ea} *lying away from the (bi-Lipschitz) horospherical boundary of* $\widetilde{M}$. *Again, let* β^h_{adm} *be the admissible quasigeodesic corresponding to* β^h_{ea} *and let* $\beta_{adm} = \beta^h_{adm} \setminus \partial\widetilde{M}$ *be the part of* β^h_{adm} *lying away from the (bi-Lipschitz) horospherical boundary of* $\widetilde{M}$.

Then $d_M(\beta_{ea}, o) \geq n$ *implies that* $d_M(\beta_{adm}, o) \geq \Theta(n)$. *Conversely,* $d_M(\beta_{adm}, o) \geq n$ *implies that* $d_M(\beta_{ea}, o) \geq \Theta(n)$.

3.5. **Cannon-Thurston Maps for Free Groups.** We identify H with its bi-Lipschitz image in M under the bi-Lipschitz homeomorphism $F : N \to M$ from the hyperbolic manifold N to the model manifold M.

We now want to show that if $\lambda = [a, b]$ is a geodesic in the intrinsic metric on $\widetilde{H}$ joining $a, b \in \widetilde{H}$, and lying outside a large ball about a fixed reference point $p \in \widetilde{H} \subset \widetilde{M}$, then the (bi-Lipschitz) hyperbolic geodesic λ_h joining $a, b \in \widetilde{M}$ also lies outside a large ball about p in $\widetilde{M}$. This would guarantee the existence of a Cannon-Thurston Map by Lemma 1.10.

To fix notation, let F be a free Kleinian group without parabolics. Let $M = \mathbb{H}^3/G$ and H a compact (Scott) core of M. $\widetilde{H}$ with its intrinsic metric is quasi-isometric to the Cayley graph Γ_F and so its intrinsic boundary may be identified with the Cantor set ∂F thought of as the Gromov boundary of Γ_F. Let $\widehat{H}$ and $\widehat{M}$ denote the compactifications by adjoining ∂F and the limit set Λ_F to $\widetilde{H}$ and $\widetilde{M}$ respectively.

Theorem 3.18. Cannon-Thurston for Free Groups *The inclusion* $i : \widetilde{H} \to \widetilde{M}$ *extends continuously to a map* $\hat{i} : \widehat{H} \to \widehat{M}$.

Proof. Let $p \in \widetilde{H}$ be a base-point, and $\lambda = [a, b]$ be a geodesic in the intrinsic metric on $\widetilde{H}$ and λ_h be the (bi-Lipschitz) hyperbolic geodesic joining its end-points in $\widetilde{M}$. By Lemma 1.10 it suffices to show that if λ lies outside a large ball about p in $\widetilde{H}$, then λ_h lies outside a large ball about p in $\widetilde{M}$.

Suppose that λ lies outside an n-ball about p in $\widetilde{H}$, i.e. $d_{\widetilde{H}}(\lambda, p) \geq n$. Let $\{D_i\}$ be a finite collection of compressing disks in H such that each component of $\partial H \setminus \bigcup_i \partial D_i$ is a pair of pants.

Since each (lift of) D_i separates $\widetilde{H}$ and since λ lies outside a large ball about p in $\widetilde{H}$, we conclude that there exists such a lift D lying outside an $m = m(n)$-ball about p in $\widetilde{H}$ and that λ lies in the component of $\widetilde{H} \setminus D$ not containing p, where $m(n) \to \infty$ as $n \to \infty$. Let $A = D \bigcup_i \sigma_i$ be the quasidisk containing D constructed in Section 3.3, where σ_i is a closed curve on the lift $\widetilde{\Sigma^s_i}$ of the $i-$th split surface Σ^s_i to $\widetilde{M}$. Also, let $\widetilde{\Sigma^s_i} \setminus \sigma_i = \widetilde{\Sigma^s_{i+}} \cup \widetilde{\Sigma^s_{i-}}$ where $\widetilde{\Sigma^s_{i+}}, \widetilde{\Sigma^s_{i-}}$ are the two components of $\widetilde{\Sigma^s_i} \setminus \sigma_i$. Similarly, let $\widetilde{H} \setminus D = \widetilde{H}_+ \cup \widetilde{H}_-$, where $\widetilde{\Sigma^s_{0+}} \subset \partial \widetilde{H}_+$ and $\widetilde{\Sigma^s_{0-}} \subset \partial \widetilde{H}_-$. Assume without loss of generality that $\lambda \subset \widetilde{H}_+$ and $p \in \widetilde{H}_-$. Let $\widetilde{M}_+ = \widetilde{H}_+ \cup \bigcup_i \widetilde{\Sigma^s_{i+}}$ and $\widetilde{M}_- = \widetilde{H}_+ \cup \bigcup_i \widetilde{\Sigma^s_{i-}}$. Also let $\widetilde{M}_H = \bigcup_i \widetilde{\Sigma^s_i}$ be the union of all the horizontal split surfaces lifted to $\widetilde{M}$.

Let α be an electro-ambient quasigeodesic joining the end-points of λ in $\widetilde{M}$ and β be an admissible quasigeodesic corresponding to α.

Since β is admissible, it consists of horizontal and vertical pieces. Two cases arise:
a) $\beta \cap \widetilde{M}_H \subset \widetilde{M}_+$
b) $\beta \cap \widetilde{M}_+ \cap \widetilde{M}_- \neq \emptyset$.

Roughly speaking Cases (a) and (b) correspond respectively to the cases where β does not or does intersect A coarsely.

Case a: $\beta \cap \widetilde{M}_H \subset \widetilde{M}_+$
By Corollary 3.14 and Lemma 3.17 it suffices to show that there exists a function $\Theta : \mathbb{N} \to \mathbb{N}$ satisfying $\Theta(n) \to \infty$ as $n \to \infty$ such that the following holds.
$d_{\widetilde{H}}(\lambda, p) \geq n$ implies that $d_M(\beta_{adm}, p) \geq \Theta(n)$.

The existence of such a function $\Theta : \mathbb{N} \to \mathbb{N}$ follows exactly as in Lemma 3.12.

Case b: $\beta \cap \widetilde{M}_+ \cap \widetilde{M}_- \neq \emptyset$
We shall say that β crosses A at x if either $x \in \beta \cap A$, or if there exists a vertical elementary admissible subpath $x \times I \subset \beta$, such that either $(x, 0) \in \widetilde{M}_+$ and $(x, 1) \in \widetilde{M}_-$ or $(x, 1) \in \widetilde{M}_+$ and $(x, 0) \in \widetilde{M}_-$.

Let r, q be the first and last points at which β crosses A. Let β_{ar}, β_{qb} be the subpaths of β joining a, r and b, q respectively. Then again, as in Case (a) above, there exists a function $\Theta : \mathbb{N} \to \mathbb{N}$ satisfying $\Theta(n) \to \infty$ as $n \to \infty$ such that

$d_{\widetilde{H}}(\lambda, p) \geq n$ implies that $d_M(\beta_{ar} \cup \beta_{qb}, p) \geq \Theta(n)$. Hence by Corollary 3.14 and Lemma 3.17 it follows that there exists a function $\Theta_2 : \mathbb{N} \to \mathbb{N}$ satisfying $\Theta_2(n) \to \infty$ as $n \to \infty$ such that $d_{\widetilde{H}}(\lambda, p) \geq n$ implies that $d_M(\mu_{ar} \cup \mu_{qb}, p) \geq \Theta_2(n)$, where μ_{ar} (resp. μ_{qb}) are the (bi-Lipschitz) hyperbolic geodesics in $\widetilde{M}$ joining a, r and b, q respectively.

If $r, q \in A$, then by Corollary 3.15, there exists a function $\Theta_3 : \mathbb{N} \to \mathbb{N}$ satisfying $\Theta_3(n) \to \infty$ as $n \to \infty$ such that $d_{\widetilde{H}}(\lambda, p) \geq n$ implies that $d_M(\mu_{rq}, p) \geq \Theta_3(n)$, where μ_{rq} is the (bi-Lipschitz) hyperbolic geodesic in $\widetilde{M}$ joining r and q. Hence by $\delta-$hyperbolicity of $\widetilde{M}$, $d_M(\mu_{ab}, p) \geq min(\Theta_2(n), \Theta_3(n)) - 2\delta$ and we are done.

Else, let $r \in \widetilde{\Sigma_i^s}$ and $q \in \widetilde{\Sigma_j^s}$. There exist $C(m), m \in \mathbb{N}$ such that $d_M(r, A) \leq C(i)$ and $d_M(q, A) \leq C(j)$. Choose r_1, q_1 in σ_i, σ_j respectively such that $d_M(r, r_1) \leq C(i)$ and $d_M(q, q_1) \leq C(j)$.

Then, again as in the proof of Lemma 3.12, there exists a function $\Theta_4 : \mathbb{N} \to \mathbb{N}$ satisfying $\Theta_4(n) \to \infty$ as $n \to \infty$ such that $d_M(\mu_{r_1 q_1}, p) \geq \Theta_4(n)$, where $\mu_{r_1 q_1}$ is the (bi-Lipschitz) hyperbolic geodesic in $\widetilde{M}$ joining r_1 and q_1. Hence by $\delta-$hyperbolicity of $\widetilde{M}$ again, $d_M(\mu_{ab}, p) \geq min(\Theta_2(n), \Theta_4(n)) - 2\delta$ and we are through. $\square$

3.6. **Finitely Generated Kleinian Groups.** In this subsection, we indicate the modifications necessary in the previous subsection to prove the analogous theorem for finitely generated Kleinian groups.

Let N_{gf} denote the augmented Scott core of $N^h = \mathbf{H}^3/G$. Let $i : N_{gf} \to N^h$ be the natural inclusion map. N_{gf} can be naturally identified with the convex core of a geometrically finite manifold including into N^h as a type-preserving homotopy equivalence. Let H be N_{gf} with open neighborhoods of cusps removed. Let N be N^h with open neighborhoods of cusps removed. Then $\widetilde{H}$ is strongly hyperbolic relative to its horospheres and $\widetilde{N}$ is strongly hyperbolic relative to its horospheres. Let $\widehat{H}$ and $\widehat{N}$ denote their relative hyperbolic compactifications. Note that $\widehat{H} = \widehat{N_{gf}}$, where $\widehat{N_{gf}}$ is the Gromov compactification of the hyperbolic space $\widetilde{N_{gf}}$. Similarly, $\widehat{N} = \widehat{N^h}$, where $\widehat{N^h}$ is the Gromov compactification of the hyperbolic space $\widetilde{N^h}$. Let $\widetilde{i} : \widetilde{H} \to \widetilde{N}$ indicate the lift of i. Let M denote the model manifold for N and let $\widehat{M}$ denote the relative hyperbolic compactification of $\widetilde{M}$. We identify H with its bi-Lipschitz image in M under the bi-Lipschitz homeomorphism $F : N \to M$. Let d_G be the graph metric on $\widetilde{M}$ equipped with a split geometry structure where all split components are incompressible.

First, suppose that H has incompressible boundary as a pared manifold. Then Theorem 1.13 shows that a Cannon-Thurston map exists for $\widetilde{i} : \widetilde{H} \to \widetilde{M}$. The point pre-image description is also furnished by Theorem 1.13.

Else H may be decomposed as the disk-connected sum of H_1, $H_2 \cdots H_{m+1}$ for manifolds H_i where at most one of the H_i's is a handlebody without any parabolics (taken to be H_1 without loss of generality) and the rest are pared manifolds with incompressible boundary. If one of the H_i's (say H_1) is a handlebody, then we choose a maximal collection of disjoint non-separating compressing disks in H_1 whose complement in H_1 is a ball. Let $D_i, i = 1 \cdots m$ denote all the compressing disks thus obtained. Since $m \geq 1$, there is at least one compressing disk.

Theorem 3.19. Cannon-Thurston for Kleinian Groups *Let* H, M, N, N_{gf}, N^h *be as above. The inclusion* $\widetilde{i} : \widetilde{H} \to \widetilde{M}$ *extends continuously to a map* $\hat{i} : \widehat{H} \to \widehat{M}$

between the relative hyperbolic compactifications. Equivalently, $\widetilde{i} : \widetilde{N_{gf}} \to \widetilde{N^h}$ *extends continuously to a map* $\hat{i} : \widehat{N_{gf}} \to \widehat{N^h}$ *between the hyperbolic compactifications.*

Proof: From Propositions 3.4, 3.7 and 3.11, we can construct quasidisks A_i corresponding to D_i as before and lift them to $\widetilde{M}$ (after partially electrocuting $\mathbb{Z}$-cusps if any).

Now, let λ be a geodesic segment in $\widetilde{H}$ lying outside a large ball $B_N(p)$ for a fixed reference point p. λ may be decomposed into (at most) three pieces λ_-, λ_0 and λ_+ as follows.

1. The middle piece λ_0 does not intersect any of the (lifts of the) compressing disks D_i in the interior. (We thus allow for the cases where λ_- and/or λ_+ are empty.)
2. the common end-point $\lambda_- \cap \lambda_0$ lies on some D_i. The same is demanded of $\lambda_0 \cap \lambda_+$.
3. the point q on λ nearest to p lies on λ_0

Two cases arise.

Case A: For a sequence of λ's lying outside larger and larger balls $B_n(p)$ about p, the last disk D_i that $[p, q]$ intersects lies outside large balls $B_m(p)$ where $m \to \infty$ as $n \to \infty$. This is exactly the case as in the proof of Theorem 3.18. The same proof goes through by Corollary 3.15.

Case B: There is (up to subsequencing) a fixed disk D_i that is the last disk that $[p, q]$ intersects. Since D_i is of uniformly bounded diameter, we may shift our base point to a point p' in the component $\widetilde{H_i}$ which is the lift of H_i 'on the other side of' D_i, i.e. having D_i on its boundary but not containing p. In this case, there exists a fixed n_0 such that λ lies outside $B_{(n-n_0)}(p')$. By shifting origin, we rewrite p' as p and $(n - n_0)$ as n.

Step 1: Now, $\lambda_0 \subset \widetilde{H_i} \subset \widetilde{H}$ as it does not meet any disk D_i in its interior. Since H_i is either a handlebody without parabolics, or a pared manifold with incompressible boundary, then by Theorem 3.18 or Theorem 1.13 respectively, a Cannon-Thurston map exists for the inclusion $\widetilde{H_i} \subset \widetilde{M}$. By (the necessity part of) Lemma 1.10 and Lemma 1.11, it follows that the hyperbolic geodesic λ_{0h} joining the end-points of λ_0 in $\widetilde{N^h}$ lies outside a large ball about p. Thus, there exists $m_1(n) \to \infty$ as $n \to \infty$ such that λ_{0h} lies outside a ball of radius $m_1(n)$ about p in $\widetilde{N^h}$.

Step 2: If λ_+ (or λ_-) is non-empty, then λ_- (or λ_+) is separated from p by a disk $D_i \subset \widetilde{H}$ lying outside $B_n(p)$.

Recall that M is a bi-Lipschitz model for N, which in trun is N^h with cusps removed. Also recall that d_G is the graph metric on $\widetilde{M}$. Then the quasidisk A_i is quasiconvex in $(\widetilde{M}, d_G)$ and lies outside a large ball of radius $m_2(n)$ about p, where $m_2(n) \to \infty$ as $n \to \infty$. Again, by constructing admissible paths and electro-ambient quasigeodesics as in the proof of Theorem 3.18, we obtain a new function $m_3(n)$ such that $m_3(n) \to \infty$ as $n \to \infty$ and so that the hyperbolic geodesics λ_{-h} or λ_{+h} lie outside a ball of radius $m_3(N)$ about p in $\widetilde{N^h}$.

Step 3: Therefore $\lambda_{-h} \cup \lambda_{0h} \cup \lambda_{+h}$ lies outside a ball of radius $m_4(n) = min\{m_1(n), m_3(n)\}$. Finally, since $\widetilde{N^h}$ is hyperbolic, the hyperbolic geodesic λ_h joining the end-points of λ lies outside a ball of radius $m(n) = m_4(n) - 2\delta$ about p. Also, $m(n) \to \infty$ as

$n \to \infty$. Therefore, by Lemma 1.10 or 1.11, it follows that the inclusion $\widetilde{i} : \widetilde{H} \to \widetilde{M}$ extends continuously to a map $\hat{i} : \widehat{H} \to \widehat{M}$ or equivalently that $\widetilde{i} : \widetilde{N_{gf}} \to \widetilde{N^h}$ extends continuously to a map $\hat{i} : \widehat{N_{gf}} \to \widehat{N^h}$. This concludes the proof. $\Box$

Let $\widehat{G_F}$ denote the Floyd compactification of a group G (See [13]). McMullen conjectured in [22] that there exists a continuous extension of $i : \Gamma_G \to \widetilde{M}$ to a map from $\widehat{G_F}$ to $\widehat{M}$. It was shown by Floyd in [13] that there is a continuous map from $\widehat{G_F}$ to $\widehat{H}$. Combining this with Theorem 3.19 above for Kleinian groups with parabolics, we get a proof of the following.

Theorem 3.20. *For any finitely generated Kleinian group G and $M = \mathbf{H}^3/G$, there is a continuous extension $\hat{i} : \widehat{G_F} \to \widehat{M}$.*

4. Point pre-images of the Cannon-Thurston Map

In this section, we determine the pre-images of points under the Cannon-Thurston map for degenerate free Kleinian groups G. The results are extended to arbitrary finitely generated Kleinian groups. We shall not have need to distinguish between the hyperbolic manifold and its bi-Lipschitz model any longer and will denote the manifold by M.

We set up some notation for the purposes of this section. Let G be a free degenerate Kleinian group without parabolics. Suppose that G is not geometrically finite. Let $M = \mathbb{H}^3/G$ be the quotient manifold. Note that the limit set of G is all of the sphere at infinity. Hence M is its own convex core. Let H be a compact core of M. H is a handlebody whose inclusion into M induces a homotopy equivalence. In fact, M deformation retracts onto H. Then $\widetilde{H}$ is embedded in $\widetilde{M} = \mathbb{H}^3$. Let Γ denote the Cayley graph of G with respect to some finite generating set of G. Assume that Γ is embedded in $\widetilde{H}$. Let S denote the boundary surface of H. We assume that the ending lamination Λ_{EL} is a geodesic lamination on S equipped with some (any) hyperbolic metric. This is well-defined up to Dehn twists along simple closed curves in S that bound disks in H. To avoid this ambiguity we will refer to the ending lamination in the Masur domain as Λ_{ELH}. $M \setminus Int(H)$ is homeomorphic to $S \times [0, \infty)$ and is bi-Lipschitz homeomorphic to an end M_S of a simply degenerate hyperbolic manifold without accidental parabolics [4] [5]. Thus $S \times [0, \infty) \subset N$ equipped with its *intrinsic* path metric is bi-lipschitz homeomorphic to M_S. We shall have need to pass interchangeably between these two below.

4.1. **EL leaves are CT leaves.** Let $i : \widetilde{H} \to \widetilde{M}$ denote the inclusion. Let ∂i denote the continuous extension of i to the boundary in Theorem 3.18. Note that the inclusion of Γ into $\widetilde{H}$ with its intrinsic metric is a quasi-isometry. So we might as well replace the inclusion of Γ into $\widetilde{N}$ by that of $\widetilde{H}$ into $\widetilde{N}$. We shall show that point pre-images under ∂i correspond to end-points of leaves of an ending lamination in the Masur domain.

The inclusion of S into H as its boundary induces a surjection of fundamental groups with infinitely generated kernel N. Let S_N denote the cover of S corresponding to N. Then $S_N \subset \widetilde{H} \subset \widetilde{M}$.

To distinguish between the ending lamination Λ_{ELH} (in the Masur domain) and bi-infinite geodesics whose end-points are identified by ∂i, we make the following definition.

Definition 4.1. *A* **CT leaf** λ_{CT} *is a bi-infinite geodesic whose end-points are identified by* ∂i.
An **EL leaf** λ_{EL} *is a bi-infinite geodesic whose end-points are ideal boundary points of either a leaf of the ending lamination, or a complementary ideal polygon.*

We shall show that

- **An** *EL leaf* **is an** *CT leaf***.**
- **A** *CT leaf* **is an** *EL leaf***.**

Proposition 4.2. EL is CT *Let* G *be a free degenerate Kleinian group without parabolics. Let* u, v *be either ideal end-points of a leaf of an ending lamination of* G*, or ideal boundary points of a complementary ideal polygon. Then* $\partial i(u) = \partial i(v)$.

Proof: This is almost identical to Proposition 2.1 of [29]. However, since the setup is somewhat different we include a proof. Take a sequence of short geodesics $\underline{s_i}$ exiting the end. Let $\underline{a_i}$ be geodesics in the intrinsic metric on the boundary S (of $\underline{H}$) freely homotopic to $\underline{s_i}$. By topological tameness [1] [14] and geometric tameness ([37] Ch. 9) we may assume further that $\underline{a_i}$'s are simple closed curves on S. Join $\underline{a_i}$ to $\underline{s_i}$ by the shortest geodesic $\underline{t_i}$ in $S \times [0, \infty)$ connecting the two curves. Then the collection $\underline{a_i}$ may be chosen to converge to the ending lamination on S ([37] Ch. 9). Also, in $S_N \subset \widetilde{H} \subset \widetilde{M}$, we obtain segments $a_i \subset \widetilde{S}$ which are finite segments whose end-points are identified by the covering map $P : \widetilde{S \times [0, \infty)} \to S \times [0, \infty)$. We also assume that P is injective restricted to the interior of a_i's mapping to $\underline{a_i}$. Similarly there exist segments $s_i \subset \widetilde{M}$ which are finite segments whose end-points are identified by the covering map $P : \widetilde{M} \to M$. We also assume that P is injective restricted to the interior of s_i's. The finite segments s_i and a_i are chosen in such a way that there exist lifts t_{1i}, t_{2i}, joining end-points of a_i to corresponding end-points of s_i. The union of these four pieces looks like a trapezium (see below, where we have omitted subscripts for convenience).



Figure: *Trapezium*

Next, given any leaf λ of the ending lamination, we may choose translates of the finite segments a_i (under the action of $\pi_1(H)$) appropriately, such that they converge to λ in S_N. For each a_i, let

$$b_i = t_{1i} \circ s_i \circ t_{2i}^{-1}$$

where t_{2i}^{-1} denotes t_{2i} with orientation reversed. If the translates of a_i we are considering have end-points lying outside large balls around a fixed reference point $p \in S_N$, it is easy to check that b_i's lie outside large balls about p in $\widetilde{M}$. Since $\mathbb{H}^3$

is δ-hyperbolic for some $\delta > 0$, it follows that the geodesic joining the end-points of b_i (and hence a_i which has the same end-points) lies in a $2\delta-$ neighborhood of b_i.

At this stage we invoke the existence theorem for Cannon-Thurston maps, Theorem 3.18. Since a_i's converge to λ and the hyperbolic geodesics joining the end-points of a_i exit all compact sets, it follows that $\partial i(u) = \partial i(v)$, where u, v denote the boundary points of λ. The Proposition follows. $\Box$

Any finitely generated Kleinian group is geometrically tame ([1] [14] [37] Ch. 9) and has finitely many ends. Observe that the proof of the above Proposition used the freeness of G only at the stage of applying Theorem 3.18. The same proof goes through verbatim for freely decomposable Kleinian groups with degenerate ends. The only modification to the above proof is that we consider one end of the manifold M at a time (and the pigeon-hole principle) along with Theorem 3.19 in place of Theorem 3.18 to obtain the following Proposition.

Proposition 4.3. **EL is CT - General Case** *Let G be a finitely generated freely decomposable Kleinian group. Let u, v be either ideal end-points of a leaf of an ending lamination of G, or ideal boundary points of a complementary ideal polygon. Then $\partial i(u) = \partial i(v)$.*

4.2. **CT leaves are EL leaves.** As usual we deal first with free degenerate groups without parabolics. We restate Theorem 1.13 in a form that we shall use. Recall that $M \setminus Int(H)$ is homeomorphic to $S \times [0, \infty)$ and is bi-Lipschitz homeomorphic to an end M_S of a simply degenerate hyperbolic manifold without accidental parabolics [4] [5]. Hence by Theorem 1.13 we have the following.

Theorem 4.4. [29] *Let S, M_S be as above. Then the inclusion $\tilde{j} : \widetilde{S} \to \widetilde{M_S}$ extends continuously to the boundary. Further, pre-images of points on the boundary are precisely ideal boundary points of a leaf of the ending lamination Λ_{ELS} of M_S, or ideal boundary points of a complementary ideal polygon whenever the Cannon-Thurston map is not one-to-one.*

We identify the Cayley graph Γ of the free group with a subset of $\widetilde{H} \subset \widetilde{M}$, viz. the orbit of a base-point joined by edges. The next Thorem is one of the main Theorems of this paper.

Theorem 4.5. *Let G be a free degenerate free Kleinian group without parabolics. Let $i : \Gamma_G \to \mathbb{H}^3$ be the natural identification of a Cayley graph of G with the orbit of a point in $\mathbb{H}^3$. Then i extends continuously to a map $\hat{i} : \widehat{\Gamma_G} \to \mathbb{D}^3$, where $\widehat{\Gamma_G}$ denotes the (Gromov) hyperbolic compactification of Γ_G. Let ∂i denote the restriction of $\hat{i}$ to the boundary $\partial \Gamma_G$ of Γ_G.*

Then $\partial i(a) = \partial i(b)$ for $a \neq b \in \partial \Gamma$ iff a, b are either ideal end-points of a leaf of an ending lamination of G, or ideal boundary points of a complementary ideal polygon.

Proof. By Theorem 3.18 the inclusion $i : \Gamma \to \widetilde{M}$ extends continuously to a map between the Gromov compactifications $\hat{i} : \widehat{\Gamma} \to \mathbb{D}^3$. Let ∂i denote the values of the above continuous extension to the boundary. Suppose $\partial i(a) = \partial i(b)$. Λ_{EL} is the ending lamination of M regarded as a subset of S. Let Λ_{ELG} denote Λ_{EL} lifted to $S_G = \partial \widetilde{H}$, which is a cover of S. We want to show that a, b are the end-points of a leaf of Λ_{ELG}. Suppose $(a, b)_\Gamma$ is the bi-infinite geodesic from a to b in $\Gamma \subset \widetilde{M}$. Assume without loss of generality that (a, b) passes through $1 \in \Gamma$. Let $a_k \to a$

and $b_k \to b$. Let $\overline{a_k b_k}$ denote the geodesic in $\widetilde{M}$ joining a_k, b_k. By continuity of the Cannon-Thurston map (Theorem 3.18) there exists $N(k) \to \infty$ as $k \to \infty$ such that $\overline{a_k b_k}$ lies outside an $N(k)$ ball about $1 \in \Gamma \subset \widetilde{M}$, where radius is measured in the hyperbolic metric on $\widetilde{M}$. Isotoping $\overline{a_k b_k}$ slightly, we can assume without loss of generality that it meets $\Gamma \subset \widetilde{M}$ only at its end-points (since Γ is one dimensional). We can further isotope $\overline{a_k b_k}$ rel. endpoints by a bounded amount (depending on the Hausdorff distance between $\widetilde{H}$ and $\Gamma \subset \widetilde{H}$) such that $\overline{a_k b_k} \cap S_G = \{c_k, d_k\}$.
1) there exist $c_k, d_k \in S_G = \partial \widetilde{H}$ with $d(a_k, c_k)$ and $d(b_k, d_k)$ uniformly bounded (independent of k)
2) if $\overline{c_k d_k}$ denotes the subpath of $\overline{a_k b_k}$ between c_k, d_k then (modifying $N(k)$ by an additive constant if necessary) $\overline{c_k d_k}$ lies outside an $N(k)$ ball about $1 \in \Gamma \subset \widetilde{M}$.
3) $\overline{c_k d_k}$ intersects $\widetilde{H}$ only at the endpoints c_k, d_k.

Thus $\overline{a_k b_k}$ is a concatenation of three pieces, $\overline{a_k c_k}$, $\overline{c_k d_k}$, $\overline{d_k b_k}$, where $\overline{a_k c_k}$ and $\overline{d_k b_k}$ are uniformly bounded in length and lie in $\widetilde{H}$, whereas $\overline{c_k d_k}$ lies in $\widetilde{M} \setminus Int(\widetilde{H})$.

Let $[c_k, d_k]_{S_G}$ denote the geodesic in the intrinsic metric on S_G which is homotopic (rel. endpoints) to $\overline{c_k d_k}$ in $\widetilde{M} \setminus Int(\widetilde{H})$. Since G is free, we can assume that its Cayley graph is a tree and (since $\widetilde{H}$ is quasi-isometric to Γ) $[c_k, d_k]_{S_G}$ passes through a point $o_k \in S_G$ at a uniformly bounded neighborhood of 1. Lift $[c_k, d_k]_{S_G}$ to some geodesic $[c_k, d_k] \subset \widetilde{S} \subset \widetilde{M_S}$ in the intrinsic metric on $\widetilde{S}$. Further assume that there exists some fixed $o \in \widetilde{S}$ such that the corresponding lift o_k' of o_k lies in a uniformly bounded neighborhood of o. Let $(\overline{c_k d_k})_{S_G}$ denote the corresponding lift of $\overline{c_k d_k}$ having the same endpoints as $[c_k, d_k]_{S_G}$ (such a choice is possible as $[c_k, d_k]_{S_G}$ and $\overline{c_k d_k}$ are homotopic rel. endpoints in the complement of $Int(\widetilde{H})$ in $\widetilde{M}$). It follows that $(\overline{c_k d_k})_{S_G}$ lies outside an $N(k)$-ball about o_k' in $\widetilde{M_S}$. Hence (modifying $N(k)$ by a further additive constant if necessary), $(\overline{c_k d_k})_{S_G}$ lies outside an $N(k)$-ball about $o \in \widetilde{M_S}$. Therefore, by the existence of Cannon-Thurston maps for $j : \widetilde{S} \to \widetilde{M_S}$ (Theorem 4.4) it follows that if $[c_\infty d_\infty]_{S_G}$ denotes any subsequential limit of the segments $[c_k, d_k]_{S_G}$ on $\widetilde{S}$, then $\partial j(c_\infty) = \partial j(d_\infty)$ and hence again by Theorem 4.4 c_∞, d_∞ are end-points of leaves of the ending lamination Λ_{ELS} of $S \subset M_S$. Finally, since (c_∞, d_∞) are bi-infinite geodesics passing through a bounded neighborhood of o, they project to leaves of Λ_{ELG} in S_G. These leaves are also well-defined as leaves of the ending lamination Λ_{ELH} as leaves of the ending lamination of M regarded as an element of the Masur domain. We have thus finally shown that $\Lambda_{CT} \subset \Lambda_{ELH}$. Combining this with Proposition 4.2 and Theorem 3.18 we have the Theorem. $\Box$

Note that in the proof of Theorem 4.5 we have used freeness of G to conclude only two things:
1) The manifold M has exactly one end.
2) The path λ in $\widetilde{H}$ can be isotoped off the Cayley graph of G embedded in $\widetilde{H}$.

To prove an analogue of Theorem 4.5 for arbitrary finitely generated Kleinian groups we continue with the notation that M is a hyperbolic manifold with augmented Scott core H. Then M has finitely many ends. We first note, that if $\lambda = (a_\infty, b_\infty)$ is a CT leaf then there exist $a_n \to a_\infty$ and $b_n \to b_\infty$ such that the geodesic realizations μ_n of $[a_n, b_n]$ in $\widetilde{M}$ leave arbitrarily large compact sets. We

may assume that $\widetilde{M} \setminus \widetilde{H}$ consists of lifts of the ends of M to $\widetilde{M}$. If μ_n intersects more than one such lift, it follows that there will be a subsegment ν_n of μ_n such that
1) ν_n is contained entirely in one of these lifts of the ends
2) Endpoints c_n, d_n of ν_n lie on $\widetilde{H}$
3) $c_n \to a_\infty$ and $d_n \to b_\infty$
We may therefore assume without loss of generality that μ_n lies in precisely one of the lifts of the ends E of M. If $S^h = H \cap E$ be its boundary then the ending lamination lies in the boundary of the (relatively) hyperbolic group $j_*(\pi_1(S^h))$ (hyperbolic relative to the cusp groups if any), where $j : S^h \to M$ is inclusion.

Fact (2) now goes through for arbitrary finitely generated Kleinian groups, as the inclusion of the augmented Scott core into M is a homotopy equivalence (in fact a deformation retract) and we are only interested in leaves which are limits of segments whose geodesic realizations lie inside the lift of a fixed end.

With this modification, and with Theorem 3.19 in place, the proof of Theorem 4.5 goes through for arbitrary finitely generated Kleinian groups. However, since we have to pick one end of the manifold at a time, the statement is a bit more involved.

Theorem 4.6. *Let G be a finitely generated Kleinian group. Let $i : \Gamma_G \to \mathbb{H}^3$ be the natural identification of a Cayley graph of G with the orbit of a point in $\mathbb{H}^3$. Then i extends continuously to a map $\hat{i} : \widehat{\Gamma_G} \to \mathbb{D}^3$, where $\widehat{\Gamma_G}$ denotes the (relative) hyperbolic compactification of Γ_G. Let ∂i denote the restriction of $\hat{i}$ to the boundary $\partial\Gamma_G$ of Γ_G.*

Let E be a degenerate end of $N^h = \mathbb{H}^3/G$ and $\widetilde{E}$ a lift of E to $\widetilde{N^h}$ and let M_{gf} be an augmented Scott core of N^h. Then the ending lamination $\mathcal{L}_E$ for the end E lifts to a lamination on $\widetilde{M_{gf}} \cap \widetilde{E}$. Each such lift $\mathcal{L}$ of the ending lamination of a degenerate end defines a relation $\mathcal{R}_\mathcal{L}$ on the (Gromov) Hyperbolic boundary $\partial\widetilde{M_{gf}}$ (equal to the relative hyperbolic boundary $\partial\Gamma_G$ of Γ_G), given by $a\mathcal{R}_\mathcal{L}b$ iff a, b are end-points of a leaf of $\mathcal{L}$. Let $\{\mathcal{R}_i\}_i$ be the entire collection of relations on $\partial\widetilde{M_{gf}}$ obtained this way. Let $\mathcal{R}$ be the transitive closure of the union $\bigcup_i \mathcal{R}_i$. Then $\partial i(a) = \partial i(b)$ iff $a\mathcal{R}b$.

5. Applications and Extensions

In this section we shall first mention a couple of applications of the main Theorems of this paper. Finally we indicate an extension of the Sullivan-McMullen dictionary between complex dynamics and Kleinian groups.

5.1. **Primitive Stable Representations.**

In [24] Minsky introduced and studied primitive stable representations, an open set of $PSl_2(\mathbb{C})$ characters of a nonabelian free group, on which the action of the outer automorphism group is properly discontinuous, and which is strictly larger than the set of discrete, faithful convex-cocompact (i.e. Schottky) characters.

In [24] Minsky also conjectured that
A discrete faithful representation of F is primitive-stable if and only if every component of the ending lamination is blocking.

Using the structure of the Cannon-Thurston map for handlebody groups, Jeon and Kim [17] have made considerable progress towards this conjecture. The connection between the Cannon-Thurston map for handlebody groups and primitive stable representations was indicated to the author by Yair Minsky in a personal communication.

We briefly sketch Jeon and Kim's argument for a degenerate free Kleinian group without parabolics. Let F be a free group of rank n. An element of F is called primitive if it is an element of a free generating set. Let $\mathcal{P}$ denote the set of bi-infinite paths w^* obtained by iterating cyclically reduced primitive elements. A representation$\rho : F \to PSl_2(\mathbb{C})$ is primitive stable if it maps elements of $\mathcal{P}$ to uniform quasigeodesics in $\mathbb{H}^3$.

Let $\{D_1, \cdots, D_k\} = \Delta$ be a system of disjoint non-separating compressing disks on a handlebody H cutting H into a 3-ball. A free generating set of F is dual to such a system. For a lamination Λ, the Whitehead graph $Wh(\Lambda, \Delta)$ is defined as follows. Cut H along Δ to obtain a planar surface
$partialH$ with $2n$ boundary components, each labeled by D_i^+ or D_i^- . The vertices of $Wh(\Lambda, \Delta)$ are the boundary circles of ∂H, with an edge whenever two circles are joined by an arc of $\Lambda \setminus \Delta$. Jeon-Kim [17] show following [24] that for the ending lamination Λ_E of a degenerate free group without parabolics, $Wh(\Lambda_E, \Delta)$ is connected and has no cutpoints.

Let ρ_E be the associated representation. If ρ is not primitive stable, then there exists a sequence of primitive cyclically reduced elements w_n such that $\rho(w_n^*)$ is not an $n-$ quasi-geodesic. After passing to a subsequence, w_n and hence w_n^* converges to a bi-infinite geodesic w_∞ in the Cayley graph with two distinct end points w_+, w_- in the Gromov boundary of F. The Cannon-Thurston map identifies w_+, w_- . Hence by Theorem 4.5 they are either the end points of a leaf of the ending lamination or ideal points of a complementary ideal polygon. Hence $Wh(w_\infty, \Delta)$ is connected and has no cutpoints. It follows that $Wh(w_n, \Delta)$ is connected and has no cutpoints for sufficiently large n. It follows from a Lemma due to Whitehead that w_n cannot be primitive for large n, a contradiction.

5.2. **Discreteness of Commensurators.** In [19] and [31], the main results of this paper are used to prove that commensurators of finitely generated, infinite covolume, Zariski dense Kleinian groups are discrete. The proof proceeds by showing that commensurators preserve the structure of point pre-images of Cannon-Thurston maps.

5.3. **Extending the Sullivan-McMullen Dictionary.** A celebrated theorem of Yoccoz in Complex Dynamics (see Hubbard [16], or Milnor [23]) proves the local connectivity of certain Julia sets using a technique called 'puzzle pieces', which consists of a decomposition of a complex domain into pieces each of which under iteration by a quadratic map converges to a single point. The dynamical system can then be regarded as a semigroup $\mathbb{Z}_+$ of transformations acting on a complex domain.

Split components can be regarded as a 3-dimensional analogue of puzzle pieces. Let us try to justify this analogy. Suppose there is a group G acting cocompactly on $\mathbb{H}^3$. Let $H \subset G$ be a subgroup. Let G/H denote the coset space. Then what we would want as the right analogue is that if one takes a sequence of elements g_i going to infinity in the coset space, the iterates of the convex hull of the limit set of H converge to a point in the limit sphere. Thus going to infinity in the coset space G/H would be the right Kleinian groups analogue of going to infinity in the semigroup $\mathbb{Z}_+$ of transformations acting on a complex domain.

In the context of this paper, H would be the fundamental group of a split component. However, for us G is a surface Kleinian group and does not act cocompactly on $\mathbb{H}^3$. We think of the quotient space $\mathbb{H}^3/H$ as parametrizing the set of normal directions to the split component. The graph metric gives a combinatorial distance on $\mathbb{H}^3/H$ and we think of $(\mathbb{H}^3/H, d_G)$ as the analogue of the semigroup $\mathbb{Z}_+$. Thus, instead of going to infinity by iteration in the semigroup $\mathbb{Z}_+$, we go to infinity in the graph metric. Further, the analogue of the requirement that iterates go to infinity, is that the visual diameter goes to zero as we move to infinity in the graph metric. This is ensured by hyperbolic quasiconvexity, and also follows easily from **graph quasiconvexity**. Note that **graph quasiconvexity** is a statement that gives uniform shrinking of visual diameter to zero as one goes to infinity.

Thus we extend the Sullivan-McMullen dictionary (see [36], [21]) between Kleinian groups and complex dynamics by suggesting the following analogy:

(1) *Puzzle pieces* are analogous to *split components*

(2) *Convergence to a point under iteration* is analogous to *graph quasiconvexity*

One issue that gets clarified by the above analogy is a point raised by McMullen in [22]. McMullen indicates that though the Julia set $J(P_\theta)$, where

$$P_\theta(z) = e^{2\pi i\theta}z + z^2$$

need not be locally connected in general by a result of Sullivan [35], the limit sets of punctured torus groups are nevertheless locally connected. Local connectivity of Julia sets would therefore not be the right analogue of local connectivity of limit sets in this setup. Instead we look at the *techniques* for proving local connectivity of limit sets vis-a-vis the *techniques* for proving local connectivity of Julia sets. Thus, by proposing the analogy between puzzle pieces and split components as above, this issue is to an extent clarified. In short, the analogy is in the technique rather than in the result.

An analogue of the $\mathbb{Z}_+$ dynamical system may also be extracted from the split geometry model. Note that each block corresponds to a splitting of the surface group, and hence an action on a tree. As $i \to \infty$, the split blocks B_i^s and hence the induced splittings also go to infinity, converging to a **free action of the surface group on an $\mathbb{R}$-tree dual to the ending lamination**. Thus iteration of the quadratic function corresponds to taking a sequence of splittings of the surface group converging to a (particular) action on an $\mathbb{R}$-tree.

Problem: The building of the Minsky model and its bi-Lipschitz equivalence to a hyperbolic manifold [25] [6] gives rise to a speculation that there should be a purely combinatorial way of doing much of the work. Bowditch's rendering [3], [4] of the Minsky, Brock-Canary-Minsky results is a step in this direction. This paper brings out the possibility that the whole thing should be do-able purely in terms of actions on trees. Of course there is an action of the surface group on a tree dual to a pants decomposition. So we do have a starting point. However, one ought to be able to give a purely combinatorial description, *ab initio*, in terms of a sequence of actions of surface groups on trees converging to an action on an $\mathbb{R}$-tree. This would open up the possibility of extending these results (including those of this paper) to other hyperbolic groups with infinite automorphism groups, notably free groups.

References

[1] I. Agol. Tameness of hyperbolic 3-manifolds. *preprint, arXiv:math.GT/0405568*, 2004.

[2] B. H. Bowditch. Relatively hyperbolic groups. *preprint, Southampton*, 1997.

[3] B. H. Bowditch. End invariants of hyperbolic manifolds. *preprint, Southampton*, 2005.

[4] B. H. Bowditch. Model geometries for hyperbolic manifolds. *preprint, Southampton*, 2005.

[5] J. F. Brock, R. D. Canary, and Y. N. Minsky. The Classification of finitely generated Kleinian groups. *in preparation.*

[6] J. F. Brock, R. D. Canary, and Y. N. Minsky. The Classification of Kleinian surface groups II: The Ending Lamination Conjecture. *preprint*, 2004.

[7] R. D. Canary. Ends of hyperbolic 3 manifolds. *J. A. M. S.*, pages 1–35, 1993.

[8] J. Cannon and W. P. Thurston. Group Invariant Peano Curves. *preprint, Princeton*, 1985.

[9] J. Cannon and W. P. Thurston. Group Invariant Peano Curves. *Geometry and Topology vol 11*, pages 1315–1356, 2007.

[10] S. Das and M. Mj. Addendum to Ending Laminations and Cannon-Thurston Maps: Parabolics. *arXiv:1002.2090*, 2010.

[11] S. Das and M. Mj. Semiconjugacies Between Relatively Hyperbolic Boundaries . *arXiv:1007.2547*, 2010.

[12] B. Farb. Relatively hyperbolic groups. *Geom. Funct. Anal. 8*, pages 810–840, 1998.

[13] W. J. Floyd. Group Completions and Limit Sets of Kleinian Groups. *Invent. Math. vol.57*, pages 205–218, 1980.

[14] D. Gabai and D. Calegari. Shrink-wrapping and the Taming of Hyperbolic 3-manifolds. *Jour. Amer. Math. Soc. 19, no. 2, arXiv:math.GT/0407161*, pages 385–446, 2006.

[15] J. Hempel. 3 manifolds. *Princeton University Press*, 1976.

[16] J. H. Hubbard. Local Connectivity of Julia sets and Bifurcation Loci: three Theorems of J.-C. Yoccoz. *In L. R. Goldberg and A. G. Philips, Editors, Topological Methods in Modern Mathematics, Publish or Perish Inc.*, pages 467–511, 1993.

[17] W. Jeon and I. Kim. Primitive stable representations of gemetrically infinite handlebody hyperbolic 3-manifolds. *preprint, arXiv:1003.2055*, 2010.

[18] E. Klarreich. Semiconjugacies between Kleinian group actions on the Riemann sphere. *Amer. J. Math 121*, pages 1031–1078, 1999.

[19] C. J. Leininger, D. D. Long, and A. W. Reid. Commensurators of non-free finitely generated Kleinian groups. *Algebraic and Geometric Topology 11*, pages 605–624, 2011.

[20] J. D. McCarthy and A. Papadopoulos. Dynamics on Thurston's Sphere of Projective Measured Foliations. *Comment. Math. Helv. 64*, pages 133–166, 1989.

[21] C. T. McMullen. Renormalization and 3-Manifolds which Fiber over the Circle. *Princeton University Press*, 1998.

[22] C. T. McMullen. Local connectivity, Kleinian groups and geodesics on the blow-up of the torus. *Invent. math.*, 97:95–127, 2001.

[23] J. Milnor. Local Connectivity of Julia sets: Expository Lectures. *In Tan Lei, Editor, The Mandelbrot Set, Theme and Variations, CUP*, pages 67–116, 2000.

[24] Y. N. Minsky. On dynamics of $Out(F_n)$ on $PSL_2(C)$ characters. *preprint, arxiv: 0906.3491*, 2009.

[25] Y. N. Minsky. The Classification of Kleinian surface groups I: Models and bounds. *Ann. of Math. 171*, pages 1–107, 2010.

[26] M. Mitra. Cannon-Thurston Maps for Trees of Hyperbolic Metric Spaces. *Jour. Diff. Geom.48*, pages 135–164, 1998.

[27] H. Miyachi. Semiconjugacies between actions of topologically tame Kleinian groups. *preprint*, 2002.

[28] M. Mj. Cannon-Thurston Maps for Surface Groups. *submitted, arXiv:math.GT/0607509*, 2006.

[29] M. Mj. Ending Laminations and Cannon-Thurston Maps. *submitted, arXiv:math.GT/0702162*, 2007.

[30] M. Mj. Cannon-Thurston Maps, i-bounded Geometry and a Theorem of McMullen. *Actes du séminaire Théorie spectrale et géométrie, Grenoble, arXiv:math.GT/0511041*, 2011.

[31] M. Mj. On Discreteness of Commensurators. *Geometry and Topology 15*, pages 331–350, 2011.

[32] M. Mj and A. Pal. Relative Hyperbolicity, Trees of Spaces and Cannon-Thurston Maps . *Geometriae Dedicata, Volume 151, Number 1, arXiv:0708.3578*, pages 59–78, 2011.

[33] J. P. Otal. Courants geodesiques et produit libres. *These d'Etat, Universit Paris-Sud, Orsay*, 1988.

[34] J. Souto. Cannon-Thurston Maps for Thick Free Groups. *preprint*, 2006.

[35] D. Sullivan. Conformal dynamical systems. *Geometric Dynamics, LNM 1007, Springer-Verlag*, pages 725–752, 1983.

[36] D. Sullivan. Quasiconformal Homeomorphisms and Dynamics I: Solution of the Fatou-Julia problem on Wandering Domains. *Annals of Math. 122*, pages 401–418, 1985.

[37] W. P. Thurston. The Geometry and Topology of 3-Manifolds. *Princeton University Notes*, 1980.